



First Cenozoic glaciers in West Antarctica

Krzysztof BIRKENMAJER¹, Andrzej GAŹDZICKI^{2,4}, Krzysztof P. KRAJEWSKI^{3,4},
Andrzej PRZYBYCIN⁴, Andrzej SOLECKI^{4,5}, Andrzej TATUR^{4,6} and Ho Il YOON⁷

¹ *Instytut Nauk Geologicznych, Polska Akademia Nauk, ul. Senacka 1-3, 31-002 Kraków, Poland
<ndbirken@cyf-kr.edu.pl>*

² *Instytut Paleobiologii, Polska Akademia Nauk, ul. Twarda 51/55, 00-818 Warszawa, Poland
<gazdzick@twarda.pan.pl>*

³ *Instytut Nauk Geologicznych, Polska Akademia Nauk, ul. Twarda 51/55, 00-818 Warszawa, Poland
<kpkraj@twarda.pan.pl>*

⁴ *Zakład Biologii Antarktyki, ul. Ustrzycka 10/12, 02-141 Warszawa, Poland*

⁵ *Instytut Nauk Geologicznych, Uniwersytet Wrocławski, pl. Maksy Borna 9, 50-204 Wrocław,
Poland <asol@ing.uni.wroc.pl>*

⁶ *Centrum Badań Ekologicznych, Polska Akademia Nauk, ul. Konopnickiej 1, 05-092 Dziekanów
Leśny, Poland <tatura@interia.pl>*

⁷ *Korea Polar Research Institute, 1270, Sa-2-dong, Sangrokgu, Ansan 426-744, Korea
<hiyoon@kordi.re.kr>*

Abstract: One of the most significant global climatic events in the Cenozoic was the transition from greenhouse to icehouse conditions in Antarctica. Tectonic evolution of the region and gradual cooling at the end of Eocene led to the first appearance of ice sheets at the Eocene/Oligocene boundary (ca. 34 Ma). Here we report geological record of mountain glaciers that preceded major ice sheet formation in Antarctica. A terrestrial, valley-type tillite up to 65 metres thick was revealed between two basaltic lava sequences in the Eocene–Oligocene Point Thomas Formation at Hervé Cove – Breccia Crag in Admiralty Bay, King George Island, South Shetland Islands. K–Ar dating of the lavas suggests the age of the glaciation at 45–41 Ma (Middle Eocene). It is the oldest Cenozoic record of alpine glaciers in West Antarctica, providing insight into the onset of glaciation of the Antarctic Peninsula and South Shetland Islands.

Key words: Antarctica, King George Island, Eocene, mountain glaciers.

Introduction

The Antarctic Cenozoic contains an important record of deterioration of global climate. The break-up of Gondwanaland, the opening of the Tasman and Drake Passages, and the progressive isolation of Antarctica by the Antarctic Circumpolar Current led to the transition from a warm, ice-free climate in early




Fig. 1. Location of Hervé Cove and Breccia Crag in Admiralty Bay on King George Island, South Shetland Islands, West Antarctica.

Eocene to a colder climate and glacial conditions at the end of Eocene and younger epochs. The first evidence of major climate change with cooling through the late Paleogene is reported from the latest Eocene or Eocene/Oligocene boundary (ca. 34 Ma). Sedimentological (Ehrmann and Mackensen 1992; Coxall *et al.* 2005), floral (Birkenmajer and Zastawniak 1989; Francis 1999, 2000), faunal (Feldmann and Woodburne 1988; Aronson and Blake 2001; Dzik and Gaździcki 2001; Reguero *et al.* 2002; Myrcha *et al.* 2002; Kriwet and Gaździcki 2003; Gaździcki 2004), and oxygen isotopic (Mackensen and Ehrmann 1992; Kennett and Warnke 1993; Gaździcki *et al.* 1992; Salamy and Zachos 1999; Dutton *et al.* 2002; Ivany *et al.* 2004) data indicate cooling, terrestrial and marine ice sheet growth, and initiation of Cenozoic glaciation at that time interval both in East and West Antarctica (Barron *et al.* 1991; Barrett 1996, 2001; Abreu and Anderson 1998; Thomson 2004). While glacial history of Antarctica has been mostly revealed from deep sea drillings (Zachos *et al.* 2001), West Antarctica shows also a fairly well-preserved glacial record exposed on South Shetland Islands and the Antarctic Peninsula (Birkenmajer 1991; Dingle and Lavalley 1998; Dingle *et*

al. 1998). The Cenozoic strata on King George Island display a sequence of alternating glacial and interglacial events, with at least two regional ice sheet expansions during the Oligocene Polonez Glaciation (32–26 Ma) and the Miocene Melville Glaciation (23–20 Ma) (Birkenmajer 2001; Troedson and Smellie 2002; Troedson and Riding 2002).

It is unclear when the first significant ice accumulations began to develop since the postulated Eocene Kraków Glaciation (Birkenmajer *et al.* 1986) was disproved using strontium isotope stratigraphy (Dingle and Lavalle 1998; Troedson and Smellie 2002). In this paper we present new evidence of alpine glaciers that preceded major ice sheet formation in West Antarctica, providing insight into the onset of Cenozoic glacial history of the region.

Geological setting

The Eocene–Oligocene Point Thomas Formation on King George Island shows a sequence of subaerial basalt to andesitic basalt, a part of the Cenozoic




Fig. 2. Sections of the Hervé Cove tillite in volcanic sequence of the Point Thomas Formation showing detailed location of samples analyzed for the K-Ar dating. For location of the sections see Fig. 3. 1 – basalt to andesitic basalt; 2 – volcanic tuffs and breccia; 3 – diamictite.




Fig. 3. Outcrops of Hervé Cove tillite at Breccia Crag (A) and Hervé Cove (B), Ezcurra Inlet, Admiralty Bay, King George Island seen from the north. BC1 to BC4 are sections illustrated in Fig. 2. BC1-1, BC1-3, BC2-1, BC2-3, BC3-1, and BC4-3 are samples analyzed for the K-Ar dating.




Fig. 4. Outcrops of Hervé Cove tillite at section BC-2 (A) and BC-3 (B). Note the unsorted nature of the diamictite dominated by boulders and debris of predominantly local rock material.

magmatic complex of the Outer Magmatic Arc of the South Shetland Islands (Birkenmajer 2001). The sequence contains many well-developed regolithic surfaces as well as fossil soil horizons with floral remains that record a gradual climatic cooling trend following the Early Eocene Climatic Optimum (Birkenmajer and Zastawniak 1989). A tillite horizon revealed at Hervé Cove and Breccia Crag (Ezcurra Inlet in Admiralty Bay) in the Point Thomas Fm provides direct evidence for the appearance of mountain glaciers during this climatic trend (Figs 1, 2). This is a valley-type tillite consisting mainly of coarse diamictite of varying thickness (10–65 m) that covers an uneven scoured surface developed on the lower basalt sequence of the formation (Figs 2, 3). Detrital components of the diamictite represent mostly local, unworked volcanic material, though exotic boulders and debris are also noted (Fig. 4). Frost-splitting of boulders and associated clastic dykes as well as washing structures, erosional surfaces and fluvial sediments evidence seasonal cool and wet climatic conditions. Petrified *Nothofagus* wood fragments and volcanic bombs and tuffs found in the tillite suggest development of glaciers in a partly forested, active volcanic mountain range. Interfingering of diamictite layers and tuffs and lava flows of the lower basalt sequence observed at marginal parts of the tillite horizon suggests that it is close in age to its substratum. The tillite is covered by tuffs and thick lava flows of the upper basalt sequence of the Point Thomas Fm (see Figs 2, 3).

Radiometric dating

Samples of basaltic lavas directly under- and overlying the Hervé Cove tillite were taken along four sections at Hervé Cove and Breccia Crag for K-Ar dating (Figs 2, 3). The dating was performed at the Korea Polar Research Institute (Ansan). Samples BC1-1, BC2-1, and BC3-1 from basalts underlying the tillite gave ages of 38.1 (± 1.4), 45.4 (± 1.2), and 41.3 (± 1.0) Ma, respectively. However, the analysis of thin sections indicates that the sample BC1-1 that yielded the age of 38.1 Ma is strongly altered, and therefore should be excluded from consideration. Samples BC1-3, BC2-3, and BC4-3 from basalts overlying the tillite gave ages of 28.6 (± 0.9), 45.1 (± 0.9), and 30.4 (± 0.7) Ma, respectively. Thus the tillite is younger than 45–41 Ma and older than 45–29 Ma. Both age limits overlap at 45–41 Ma, and that value range is suggested as the most probable age of the Hervé Cove tillite.

Concluding remarks

Global deep-sea oxygen isotope curve suggests rapid development of Antarctic ice sheets close to the Eocene/Oligocene boundary (Zachos *et al.* 2001). This abrupt climatic event took place at the end of the 15 Ma-long cooling trend




Fig. 5. Cenozoic climatic history of West Antarctica and main glacial events revealed from sedimentary record on King George Island. Global deep-sea oxygen isotope curve and climatic trends after Zachos *et al.* (2001). Geologic time scale after Gradstein and Ogg (2004).

following the Early Eocene Climatic Optimum (Fig. 5). Gradual deterioration of climate during this time interval is recorded in both the marine and terrestrial realms. Progressive decrease of sea water temperature can be deduced from geological data (sea-level changes, deposition of dropstones in open sea sediments), from geochemical proxies (stable oxygen isotope record, clay mineral composition) as well as from paleontological investigations (dinoflagellates, diatoms, radiolarians, foraminifers and molluscs) of deep-sea cores (Kennett and Warnke 1992, 1993). Important changes of the terrestrial ecosystem resulting in progradation of *Nothofagus*-dominated forests have been noted at lower latitudes on Seymour Island (Antarctic Peninsula) as well as at higher latitudes in McMurdo Sound in Antarctica (Case 1988; Doktor *et al.* 1996; Francis 1999, 2000; Pole *et al.* 2000). It is suggested by several authors that mountain glaciers might have developed in the forested Antarctic landscape inhabited by rich vertebrate faunas of marsupials and ratite birds (Prothero 1994; Tambussi *et al.* 1994; Reguero *et al.* 2002). However, there has been no direct evidence of such glacial activity re-




Fig. 6. Eocene paleogeographic and surface circulation reconstruction of the Southern Hemisphere and the location of first alpine glaciers in West Antarctica. Paleogeography and surface circulation after Lazarus and Caulet (1993).

ported. We document it here for the first time. Mountain glaciers occurred in the marginal zone of the Antarctic continent in the Middle Eocene (Fig. 6).

Acknowledgements. — The field work in Antarctica was carried out during the 27th Polish Antarctic Expedition to *Arctowski* Station in austral summer of 2002–2003. Professor Robert H. Rutford and two anonymous referees gave helpful comments on the manuscript. This study was supported in part by grant from the Polish Committee for Scientific Research (PBZ-KBN-108/PO4/1) and Korean Polar Research Program (PE05004/PN044001/PN04002). A contribution to the SCAR Scientific Research Programme *Antarctic Climate Evolution* (ACE).

References

- ABREU V.S and ANDERSON J.B. 1998. Glacial eustasy during the Cenozoic: sequence stratigraphic implications. *American Association of Petroleum Geologists Bulletin* 82: 1385–1400.
- ARONSON B.R. and BLAKE D.B. 2001. Global climate change and origin of modern benthic communities in Antarctica. *American Zoologist* 41: 27–39.
- BARRETT P.J. 1996. Antarctic palaeoenvironment through Cenozoic time – a review. *Terra Antarctica* 3: 103–119.
- BARRETT P.J. 2001. Climate change – an Antarctic perspective. *New Zealand Science Review* 58: 18–23.

- BARRON J., LARSEN B. and BALDAUF J.G. 1991. Evidence for Late Eocene to Early Oligocene Antarctic glaciation and observations on Late Neogene glacial history of Antarctica: results from Leg 119. In: J. Barron, B. Larsen and J.G. Baldauf (eds) *Proceedings of the Ocean Drilling Program, Scientific Results* 119: 869–891.
- BIRKENMAJER K. 1991. Tertiary glaciation in the South Shetland Islands, West Antarctica: evaluation of data. In: M.R.A. Thomson, J.A. Crame and J.W. Thomson (eds) *Geological Evolution of Antarctica*. Cambridge University Press: 629–632.
- BIRKENMAJER K. 2001. Mesozoic and Cenozoic stratigraphic units in parts of the South Shetland Islands and Northern Antarctic Peninsula (as used by the Polish Antarctic Programmes). *Studia Geologica Polonica* 118: 5–188.
- BIRKENMAJER K. and ZASTAWNIAK E. 1989. Late Cretaceous–early Tertiary floras of King George Island, West Antarctica: their stratigraphic distribution and paleoclimatic significance. In: J.A. Crame (ed.) *Origins and evolution of the Antarctic biota*. Geological Society Special Publication 47: 227–240.
- BIRKENMAJER K., DELITALA M.C., NARĘBSKI W., NICOLETTI M. and PETRUCCIANI C. 1986. Geochronology of Tertiary island-arc volcanics and glaciogenic deposits, King George Island, South Shetland Islands (West Antarctica). *Bulletin Polish Academy of Sciences: Earth Sciences* 34: 257–273.
- CASE J.A. 1988. Paleogene floras from Seymour Island, Antarctic Peninsula. In: R.M. Feldmann and M.O. Woodburne (eds) *Geology and paleontology of Seymour Island, Antarctic Peninsula*. Geological Society of America, Memoir 169: 523–530.
- COXALL H.K., WILSON P.A., PÁLIKE H., LEAR C.H. and BECKMAN J. 2005. Rapid stepwise onset of Antarctic glaciation and deep calcite compensation in the Pacific Ocean. *Nature* 433: 53–57.
- DINGLE R.V. and LAVELLE M. 1998. Antarctic Peninsular cryosphere: Early Oligocene (c. 30 Ma) initiation and a revised glacial chronology. *Journal of the Geological Society London* 155: 433–437.
- DINGLE S.A., MARENSSI S.A. and LAVELLE M. 1998. High latitude Eocene climate deterioration: evidence from the northern Antarctic Peninsula. *Journal of South American Earth Sciences* 11: 571–579.
- DOKTOR M., GAŹDZICKI A., JERZMAŃSKA A., POREBSKI S.J. and ZASTAWNIAK E. 1996. A plant-and-fish assemblage from the Eocene La Meseta Formation of Seymour Island (Antarctic Peninsula) and its environmental implications. In: A. Gaździcki (ed.) *Palaeontological Results of the Polish Antarctic Expeditions. Part II. Palaeontologia Polonica* 55: 127–146.
- DUTTON A.L., LOHMANN K.C. and ZINSMEISTER W.J. 2002. Stable isotope and minor element proxies for Eocene climate of Seymour Island. *Paleoceanography* 17: 6-1–6-13.
- DZIK J. and GAŹDZICKI A. 2001. The Eocene expansion of nautilids to high latitudes. *Palaeogeography Palaeoclimatology Palaeoecology* 17: 279–312.
- EHRMANN W.U. and MACKENSEN A. 1992. Sedimentological evidence for the formation of an East Antarctic ice sheet in Eocene/Oligocene time. *Palaeogeography Palaeoclimatology Palaeoecology* 93: 85–112.
- FELDMANN R.M. and WOODBURNE M.O. (eds) 1988. *Geology and paleontology of Seymour Island, Antarctic Peninsula*. Geological Society of America, Memoir 169: 1–566.
- FRANCIS J.E. 1999. Evidence from fossil plants for Antarctic palaeoclimates over the past 100 million years. *Terra Antarctica Reports* 3: 43–52.
- FRANCIS J.E. 2000. Fossil wood from Eocene high latitude forests. In: J.D. Stilwell and R.M. Feldmann (eds) *Paleobiology and paleoenvironments of Eocene rocks, McMurdo Sound, East Antarctica*. Antarctic Research Series 76: 253–260.
- GAŹDZICKI A. 2004. Eocene–Pliocene ecosystems of South Shetlands and Antarctic Peninsula. *Bollettino di Geofisica Teorica ed Applicata* 45: 258–260.
- GAŹDZICKI A., GRUSZCZYŃSKI M., HOFFMAN A., MAŁKOWSKI K., MARENSSI S.A., HAŁAS S. and TATUR A. 1992. Stable carbon and oxygen isotope record in the Paleogene La Meseta Formation, Seymour Island, Antarctica. *Antarctic Science* 4: 461–468.

- GRADSTEIN F.M. and OGG J.G. 2004. Geologic Time Scale 2004 – why, how, and where next! *Lethaia* 37: 175–181.
- IVANY L.C., BLAKE D.B., LOHMANN K.C. and ARONSON R.B. 2004. Eocene cooling recorded in the chemistry of La Meseta Formation mollusks, Seymour Island, Antarctic Peninsula. *Bollettino di Geofisica Teorica ed Applicata* 45: 242–245.
- KENNETT J.P. and WARNKE D.A. (eds) 1992. *The Antarctic paleoenvironment: a perspective on global change. Part One*. Antarctic Research Series 56: 1–385.
- KENNETT J.P. and WARNKE D.A. (eds) 1993. *The Antarctic paleoenvironment: a perspective on global change. Part Two*. Antarctic Research Series 60: 1–273.
- KRIWET J. and GAŹDZICKI A. 2003. New Eocene Antarctic chimeroid fish (Holocephali, Chimaeriformes). *Polish Polar Research* 24 (1): 29–51.
- LAZARUS D. and CAULET J.P. 1993. Cenozoic Southern Ocean reconstructions from sedimentologic, radiolarian, and other microfossil data. In: J.P. Kennett and D.A. Warnke (eds) *The Antarctic paleoenvironment: a perspective on global change. Part Two*. Antarctic Research Series 60: 145–174.
- MACKENSEN A. and EHRMANN W.U. 1992. Middle Eocene through Early Oligocene climate history and paleoceanography in the Southern Ocean: stable oxygen and carbon isotopes from ODP sites on Maud Rise and Kerguelen Plateau. *Marine Geology* 108: 1–27.
- MYRCHA A., JADWISZCZAK P., TAMBUSI C.P., NORIEGA J.I., GAŹDZICKI A., TATUR A. and DEL VALLE R.A. 2002. Taxonomic revision of Eocene Antarctic penguins based on tarsometatarsal morphology. *Polish Polar Research* 23: 5–46.
- POLE M., HILL B. and HARWOOD D. 2000. Eocene plant macrofossils from erratics, McMurdo Sound, Antarctica. In: J.D. Stilwell and R.M. Feldmann (eds) *Paleobiology and paleoenvironments of Eocene rocks, McMurdo Sound, East Antarctica*. Antarctic Research Series 76: 243–251.
- PROTHERO D.R. 1994. The Late Eocene–Oligocene extinctions. *Annual Review of Earth and Planetary Sciences* 22: 145–165.
- REGUERO M.A., MARENSSI S.A. and SANTILLANA S.N. 2002. Antarctic Peninsula and South America (Patagonia) Paleogene terrestrial faunas and environments: biogeographic relationships. *Palaeogeography Palaeoclimatology Palaeoecology* 179: 189–210.
- SALAMY K.A. and ZACHOS J.C. 1999. Latest Eocene–Early Oligocene climate change and Southern Ocean fertility: inferences from sediment accumulation and stable isotope data. *Palaeogeography Palaeoclimatology Palaeoecology* 145: 61–77.
- TAMBUSI C.P., NORIEGA J.I., GAŹDZICKI A., TATUR A., REGUERO M.A. and VIZCAINO S.F. 1994. Ratite bird from the Paleogene La Meseta Formation, Seymour Island, Antarctica. *Polish Polar Research* 15: 15–20.
- THOMSON M.R.A. 2004. Geological and palaeoenvironmental history of the Scotia Sea region as a basis for biological interpretation. *Deep-Sea Research II* 51: 1467–1487.
- TROEDSON A.L. and RIDING J.B. 2002. Upper Oligocene to lowermost Miocene strata of King George Island, South Shetland Islands, Antarctica: stratigraphy, facies analysis and implications for the glacial history of the Antarctic Peninsula. *Journal of Sedimentary Research* 72: 510–523.
- TROEDSON A.L. and SMELLIE J.L. 2002. The Polonez Cove Formation of King George Island, Antarctica: stratigraphy, facies and implications for mid-Cenozoic cryosphere development. *Sedimentology* 49: 277–301.
- ZACHOS J., PAGANI M., SLOAN L., THOMAS E. and BILLUPS K. 2001. Trends, rhythms, and aberrations in global climate 65 Ma to Present. *Science* 292: 686–693.

Received 12 October 2004

Accepted 7 January 2005