

BIOLOGY OF *HIPPODAMIA VARIEGATA* (GOEZE) (COLEOPTERA: COCCINELLIDAE), ON *APHIS FABAE* SCOPOLI (HEMIPTERA: APHIDIDAE)

Reza Jafari*

Faculty of Agriculture, Borojerd Islamic Azad University, Borojerd, Iran

Received: February 12, 2010

Accepted: March 11, 2011

Abstract: *Hippodamia variegata* (Goeze) is a among natural enemy of bean aphid so far identified in the world. In this study, the duration of hatching, larval instars and pupal stages of *H. variegata* feeding on black bean aphid, *Aphis fabae* Scopoli were investigated under laboratory conditions (25±1°C, 60±5% relative humidity and 16:8 (light:darkness) hours photoperiod. The mean pre-oviposition period of *H. variegata* was 6.20±0.13 days and the oviposition period was 43±0.21 days. The female beetles laid on an average 943.90±53.53 eggs and the hatching % was 82.86±3.12. The average incubation period was 3.35±0.11 days and the duration of total larval period from egg to adult was 16.5±0.13 days. The pre-pupal and pupal periods were 1.40±0.16 and 3.10±0.07 days, respectively. The mean longevity of the male and the female beetles were 50±3.2 and 55.5±3.37 days, respectively.

Key words: *Hippodamia variegata*, development, *Aphis fabae*

INTRODUCTION

Aphids are a diverse group of plant feeding insects belong to family Aphididae of order Hemiptera, predominantly found in temperate climate zones such as north America, Europe, Central and Eastern Asia (Nelson and Rosenheim 2006). They cause damage directly by sucking cell sap, secrete honeydew resulting in development of sooty mould on leaves and shoots and directly as vectors of certain plant viruses (Kennedy *et al.* 1962; Raboudi *et al.* 2002). *Aphis fabae* Scopoli, commonly known as black bean aphid, is the most important polyphagous species of pest and attacks the bean plant and over hundred host plants including vegetables, fruits trees and ornamental by the nymphs and the adults of aphid cause damage by sucking the sap from the tender shoots, pods, flowers and reduce the market value (Völkl and Stechmann 1998).

In Iran the bean growers use various insecticides to control black bean aphid. Insecticidal control is not only expensive but also its residues left over the sprayed surface of the crops or in the soil and have become a matter of concern of environmental pollution. The indiscriminate use of pesticides causes phytotoxicity and destruction of beneficial organisms such as predators, parasitoids, microorganisms and pollinators (Luckman and Metcalf 1978). Global warning has cautioned us and the adverse consequences of insecticide use are always alarming and also inducing pest out break because of pest resistance. These entomological backlashes have compelled the scientists to be concerned with entomologically compatible pest management programs (Hodek 1970). Therefore, use

of natural enemies in biological control is a good pest management tactic to minimize the population level of invasive pests (Delfoss 2005). Investigation into the potential use of predatory insects or mites to control aphid pests received a greater attention for a reduction of chemical damages to the environment.

The ladybird beetles have been known worldwide as a predator of a number of insects. They are the most commonly known of all beneficial insects (William 2002). The coccinellid beetles are considered to be a great economic importance in agro-ecosystem through their successful employed in the biological control of many injurious insect (Agarwala and Dixon 1992). The degree of their adaptation as well as their efficiency in controlling aphid populations observed varies with the species and the environmental conditions (Dixon 2000). They are of great economic important as predators both in their larval and adult stages on various important crop pests such as aphids, coccids and other soft bodied insects (Hippa *et al.* 1978; Kring *et al.* 1985), while the species *Hippodamia variegata* feed on many insect pests such as aphids, psyllids, white flies, lepidopteron insects and mealy bug (Franzman 2002).

The species of the genus *Hippodamia* have been identified so far in the word are reported as Aphid predators in all agro-ecosystems. Several of them have been identified as potential biological control agents of bean aphid in agricultural crops (Fan and Zhao 1988). The voracious feeding of injurious aphids by *Hippodamia* spp. keep them as appropriate effective biological control agents (Kontadimas and Stathas 2005). The searching ability of this

*Corresponding address:

Jafari_po@yahoo.com

winged predators, enable them for better locating and identifying aphid colonies under outdoor and indoor conditions as compared to other aphids natural enemies. According to Debach and Rosen (1991) a predator is a reliable regulator of its prey at low densities, if he has a great capacity for research, which would enable it to prevent the prey of causing economic damage. A similar concept has also been demonstrated by Pedigo (2004) if a predator is an effective regulator of its prey, it is able to regulate its population at low densities and prevent inflict damage to the host plant. In this context, the voracious aphid predator *H. variegata* (palaeartic species), is reported as one of the most important bio-control agents of black bean aphid in agricultural crops and it is now released in many countries to control *Aphis fabae*.

The study of the life history and the feeding behavior of *H. variegata* under laboratory conditions may evaluate its predatory impacts on black bean aphid pest for better utilization in integrated aphid control program. The study of the biology of *H. variegata* would help to use this insect for in proper biological control. Lack of research on the biology and its beneficial utilization leads to the present work.

MATERIALS AND METHODS

Collection and mass culture of *H. variegata* and *A. fabae*

To maintain the culture of black bean aphid, sugar beet (*Beta vulgaris*) and common bean (*Phaseolus vulgaris*) were grown in the micro-plots sized 2x3 meter, plants were maintained at experimental field of the Department of Culture, Faculty of Agricultural Sciences, Boroujerd Islamic Azad University, Boroujerd, Iran.

A culture of large number of larvae and adult predator of *H. variegata* was established in the laboratory in order to supply necessary insects for the experiment. For this reason, some males and females of the *H. variegata* were collected by sweep net from the unsprayed horticulture field and were confined in cages (7x7x7 cm³). Black bean aphids were also collected daily they free on infested bean leaves, stems, twigs and inflorescences from the same unsprayed field and supplied as food. These beetles were sexed and paired in Plexiglas cage (6x11x23 cm³). The bottom of the cages was covered with blotting paper.

Study of biology

During intensive observation the leaves containing *H. variegata* eggs were collected immediately after egg laying and transferred in the Petri dishes (12 cm diameter). Egg laid by each female during 24 hours were counted and kept in separate Petri dish to determine the total number of eggs laid per female and also hatching period as well as hatching percentage of eggs were recorded. After hatching of eggs, the larvae were then transferred individually in Petri dishes (15 cm diameter) and reared on black bean aphids. Fresh bean shoots infested with aphids were placed in each Petri to provide food for the larvae every morning. The larvae were observed daily at 12 hrs interval until pupation. Side by side, the exuviae found in each of vials was removed when the larva entered into the next in-

star. Total number of aphid consumed by each larva, dead and unconsumed aphids were counted daily and were replaced by fresh aphids. They multiply parthenogenetically and their density increased rapidly on the growing points of plants. The larvae and pupae of *H. variegata* were collected from naturally infested field crops and kept in incubator for adult emergence. The condition maintained in incubator was 25±1°C temperature coupled with, 60±5% relative humidity and the photoperiod 16:8 (L : D). Freshly emerged adult beetles were reared on black bean aphid in pairs in Plexiglas (6x11x23 cm³). A blotting paper was spread over its inner surface for egg laying. Fecundity, longevity and sex ratio were recorded. The emerged adults were sexed and confined in pairs in the Petri dishes and fed on black bean aphids. Sufficient food (*A. fabae* of various stages) was supplied regularly. Egg masses were collected and reared as above and continued for several times for obtaining large number of larvae and adult predators. Regular observations were made on the leaves of aphid-infested. Ten replications were used in this case.

Statistical analysis

The Data were analyzed by Analysis of variance (ANOVA) and the mean values were separated by Duncan multiple range test (DMRT). All analysis were performed using SAS, version 7.

RESULTS AND DISCUSSION

Development

The developmental stages of *H. variegata* observed as egg, four larval instars, pupa and adult. Egg is elongate, oval and generally pale yellow color. Gradual color changes occur from yellow to orange during the incubation period. Eggs were laid either singly or in the form of patches (17–25 eggs/site). Larva is blackish in color with many long branched hairs and black patches on upper surface of the body. The larval period during I and II instars were more active with a preference to prey eggs. The same phenomenon for III and IV instars utilizing more nymphal and adult prey stages was recorded. As the larva matures, it becomes Yellowish, with black spots on the dorsal surface. All prey stages were attacked by the four larval stages as well as the adult predator. This kind of prey feeding was also reported by some researchers (Elhagh and Zaiton 1996; Kontadimas and Stathas 2005). But larval stages of the predator prefer immature stages of prey aphid whereas adult predator was mostly observed feeding either nymph or adult prey aphid. Pupa is the only inactive stage of the coccinellid with flattened blackish color at early pupal stage that turned later into orange reddish before emerging as adult stage. Medium reddish adult predator emerged from pupa and searched different sites of leaf lamina for *A. fabae*. Kontadimas and Stathas (2005), also reported similar feeding ability by *H. variegata*. The drop excretion on tip of cornicle by bean aphids was reported as a barrier of predation while these phenomena was observed without any negative effect on the rate of prey aphid feeding by *H. variegata*. This was due to the strong mandibles of the predator.

Table 1. Life cycle of *H. variegata* reared on *A. fabae*

Different stages	Duration of days		
	minimum	maximum	mean ±SE
Pre-oviposition period	6	7	6.20±0.13
Ovipoisition period	37	48	3.35±0.11
Incubation period	3	4	3.35±0.11
Total larval period	13	18.5	16.5±0.13
Pre-pupal period	1	2	1.40±0.16
Pupal period	3	3.5	3.10±0.07
Total period from egg to adult	20	23.5	21.35±0.39
Longevity of the beetle	30	70	55.50±3.37

Table 2. Larval-instar diuration of *H. variegata* reared on *Aphis fabae*

Larval-instar	Mean	±SE
First	3.5	0.17
Second	3.05	0.2
Third	3.3	0.11
Fourth	3.65	0.13




Fig. 1. Duration (mean± SE) of different larval instars of *H. variegata* reared on *A.fabae*

Oviposition

The pre-oviposition period of *H. variegata* was 6 to 7 days with an average of 6.20±0.13 days (Table 1). Lanzoni *et al.* (2004) observed that the pre-oviposition period was 7 day. Elhagh and Zaiton (1996) studied that the pre-oviposition period was 6 to 7 days, which is similar to the present findings.

The oviposition period in the present study was 37 to 48 days (Table 1), and the mean oviposition period was 43±0.21 days. Elhagh and Zaiton (1996) reported that the period of *H. variegata* lasted from 35 to 48 days using *A. fabae* as food, which is similar to the present observation.

The incubation period was 3 to 4 days with an average of 3.35±0.11 days (Table 1). Elhabi *et al.* (2000) observed that the incubation period of *H. variegata* was 3 days. However, Lanzoni *et al.* (2004) and Grigorov (1977) reported that it was 3 and 3 days respectively using bean aphid as host. These results seem to be close with the present observation.

The mean of total larval period (1st instar to 4th instar) was 16.5±0.13 days (Table 1). Kontadimas and Stathas (2005) observed that the total period of *H. variegata* was 15 days using *Dyaphis crataegi* as food, which is low-

er than the present finding. Grigorov (1977) observed that the total larval period of *H. variegata* varied from 17 to 19 days on bean aphid. This result was higher than the present study. However, Wang *et al.* (2004) reported that the quality of food and environmental factors like temperature, humidity also play an important role on different aspects of the biology of coccinellid beetles. So, this variation may be due to the quality of food and environmental factors like temperature and humidity.

The newly hatched larval period was 3 to 4 days and on an average of 3.50±0.17 days (Table 2). Lanzoni *et al.* (2004) reported that this period of *H. variegata* was 2 to 3 days using bean aphid as host. This result supported to the present observation. The result of the duration of the 2nd instar varied from 2 to 4 days and the mean duration was 3.05±0.20 days (Table 2). Lanzoni *et al.* (2004) founded that the duration of the 2nd instar larvae of *H. variegata* was 2.5 to 4 days on aphid, which is comparatively similar to the results of the present findings. Elhabi *et al.* (2000) founded that the duration of 2nd instar of *H. variegata* varied from 2.5 to 3.5 days using bean aphid as a host. The result indicated that the duration of the 3rd instar larvae lasted from 3 to 4 days. The mean

duration of 3rd instar larvae was 3.30 ± 0.11 days (Table 2). Kontadimas and Stathas (2005) founded that the duration of 3rd instar larvae of *H. variegata* varied from 2.5 to 4 days on cotton aphid. This result supported to the present observation. Observation made on the larval duration of the 4th instar larvae on an average 3.65 ± 0.13 days with minimum of 3 days and maximum of 4 days (Table 2). Lanzoni *et al.* (2004) reported that the duration of final instar larvae of *H. variegata* was 3 days. Kontadimas and Stathas (2005) found that the duration of 4th instar larvae of *H. variegata* varied from 2.8 to 3.6 days on cotton aphid. This duration was within the range of present findings.

The pupal period was 3 to 3.5 days with an average 3.10 ± 0.07 days (Table 1). Wang *et al.* (2004) recorded that the mean pupal duration of *H. variegata* was 4.15 ± 1.1 days when larvae reared on *D. noxia*. Different finding revealed that the pupal period of coccinellid beetles varied with the differences of food and it was correlated with the temperature (Wang *et al.* 2004).

The number of eggs laid per female were 587 to 1247 with an average 943.90 ± 53.53 .

The 60 to 70% of total fecundity was observed within the first 23 days of ovipositional period. The maximum daily fecundity recorded (103 eggs/day) was found correlated with the peak of predation ability. The mean hatching percentage were 82.86 ± 3.12 . Lanzoni *et al.* (2004) reported that the number of eggs deposited per female of *H. variegata* was 900 ± 80.23 and 70% eggs were hatched. Elhabib *et al.* (2000) observed that the fecundity of female varied from 800–900 eggs with mean 870.5 and with average 79% eggs were hatched. These results seem to be close with the present findings.

The longevity of adult beetles was counted from the emergence of the adult to its death. The longevity of the male beetles varied from 30 to 62 days with an average of 50 ± 3.2 days. Whereas, the longevity of the female beetles varied from 30 to 70 days with an average of 55.5 ± 3.37 days. It showed that the longevity of the male beetle was shorter than the female. But there were no significant differences. Elhag and Zaiton (1996) reported that the adult of *H. variegata* lived for 32 to 60 days. This result was near similar to the present study. Elhabib *et al.* (2000) found that the longevity of male and female were 44 ± 2 and 61 ± 9.89 days. These results are also close to the present observation.

ACKNOWLEDGEMENTS

The equipment for this study is supported by Islamic Azad University, Boroujerd Branches of Iran. We gratefully acknowledge the Management of this university for their financial and technical support.

REFERENCES

- Agarwala R.K., Dixon A.F.G. 1992. Laboratory study of cannibalism and interspecific predation in ladybirds. *Ecol. Entomol.* 17 (3): 303–330.
- Debach P., Rosen D. 1991. *Biological Control by Natural Enemies*. McMillan com. New York, London, 567.
- Delfoss E.S. 2005. Risk and ethics in biological control. *Biol. Control* 35 (2): 319–329.
- Dixon A.F.G. 2000. *Insect Predator-Prey Dynamics Lady Birds Beetles and Biological Control*. New York. Cambridge University Press, 257 pp.
- Elhag E.T.A., Zaiton A.A. 1996. Biological parameters for four coccinellid species in central Saudi Arabia. *Biol. Control* 7 (3): 316–319.
- Elhabib M., Sekkat A., Eljadd L., Boumezzough A. 2000. Biology of *Hippodamia variegata* and its suitability under greenhouse conditions. *J. Appl. Entomol.* 124 (9/10): 365–374.
- Fan G.H., Zhao J.F. 1988. Functional response of *Adonia variegata* to cotton aphids. *Nat. Econ. Insect* 10 (4): 187–190.
- Franzman B.A. 2002. *Hippodamia variegata* a predacious ladybird new in Australia. *Aust. J. Entomol.* 41 (4): 375–377.
- Grigorov S. 1977. Biological peculiarities of some ladybird beetle attacking aphids. *Oecologia* 87 (4): 401–408.
- Hippa H., Kepeken S.D., Laine T. 1978. On the feeding biology of *Coccinella hieroglyphica* L. (Coleoptera, Coccinellidae). *Kevo-subarctic Ras. Station* 14 (2): 18–20.
- Hodek I. 1970. Coccinellids and the modern pest management. *Bioscience* 20: 543–552.
- Kennedy J.S., Day M.F., Eastop V.F. 1962. *A Conspectus of Aphids as Vectors of Plant Viruses*. Commonwealth Inst. Ent., London, 114 pp.
- Kontadimas D.C., Stathas G.J. 2005. Phenology fecundity and life table parameters of the predator *Hippodamia variegata* reared on *Dyaphis crataegi*. *Biocontrol* 50 (2): 223–233.
- Kring T.J., Gilstrap F.E., Michels G.I. 1985. Role of indigenous coccinellid in regulating green bugs on Texas grain sorghum. *J. Econ. Entomol.* 78 (1): 269–273.
- Lanzoni A., Accinelli G., Bazzocchi G., Burgio G. 2004. Biological traits and life table of the exotic *Harmonia axyridis* compared with *Hippodamia variegata* and *Adalia bipunctata*. *J. Appl. Entomol.* 128 (4): 298–306.
- Luckman W.H., Metcalf R.L. 1978. The pest management concept. p. 3–35. In: "Introduction to Insect Pest Management". Willey, New York, 257 pp.
- Nelson E.H., Rosenheim J.A. 2006. Encounters between aphids and their predators: The relative frequencies of disturbance and consumption. *Entomol. Exp. Appl.* 118 (1): 211–219.
- Pedigo L.P. 2004. *Entomology and Pest Management*. Prentice-Hall of India Pvt. Ltd. New Delhi, 210 pp.
- Raboudi F., Benmoussa A., Makni H., Marrakchi M. 2002. Serological detection of plant viruses in their aphid vectors and host plants in Tunisia. *Bull. OEPP/EPPO Bull.* 32 (4): 495–498.
- Volkl W., Stechmann D.H. 1998. Parasitism of the black bean aphid (*Aphis fabae*) by *Lysiphlebus fabarum* (Hymenoptera, Aphididae): the influence of host plant and habitat. *J. Appl. Entomol.* 122: 201–206.
- Wang J., Liyon C., Yang D. 2004. A study on the hunger tolerance of *Hippodamia variegata* in north Xingjian. *J. China Cott.* 31 (5): 12–14.
- William F.L. 2002. *Lady Beetles*. Ohio State University Extension Fact Sheet, Horticulture and Crop Science. Division of Wildlife, 2021 Coffey Rd. Columbus, Ohio, 857 pp.

POLISH SUMMARY**BIOLOGIA *HIPPODAMIA VARIEGATA* GOEZE (COLEOPTERA: COCCINELLIDAE), NA *APHIS FABAE* SCOPOLI (HEMIPTERA: APHIDIDAE)**

Hippodamia variegata Goeze jest wrogiem naturalnym mszycy fasoli określonym w skali światowej. W pracy, w warunkach laboratoryjnych ($25\pm 1^{\circ}\text{C}$, 60–65% wilgotności względnej i fotoperiodzie 16 8 godzin – światłości, ciemności), badano czas trwania wylęgu, stadia larwalne

i stadia poczwarki *H. variegata*. Średni okres czasu przed złożeniem jaj *H. variegata* wynosił $6,20\pm 0,13$ dni, a okres składania jaj – $43\pm 0,21$ dni. Żeńskie osobniki chrząszcza składały średnio $943,90\pm 53,53$ jaj, a procent wylęgu wynosił $82,86\pm 3,12$. Średni czas okresu inkubacji wynosił $3,35\pm 0,11$ dni, a czas trwania całkowitego okresu larwalnego od jaja do dorosłości – $16,5\pm 13$ dni. Okresy poprzedzające stadium przedpoczwarki i poczwarki wynosiły, odpowiednio $1,40\pm 0,16$ i $3,10\pm 0,07$ dni. Średni czas życia dla chrząszczy męskich i żeńskich wynosił, odpowiednio $50\pm 3,2$ i $55,5\pm 3,37$ dni.