

Effects of heavy metals on bioremediation of diesel by Antarctic microalga *Tritostichococcus* sp.

Wen Xi GOH¹ ORCID-0009-0003-9761-397X, Ruo-Shan TAE¹ ORCID- 0009-0007-4122-4085, Zheng Syuen LIM^{1,2} ORCID- 0000-0002-8650-8408, Chiew Yen WONG^{1,3*} ORCID- 0000-0003-0534-6206, Siti Aqlima AHMAD² ORCID- 0000-0002-7625-3704, Yih Yih KOK¹ ORCID- 0000-0002-4999-7126 and Peter CONVEY^{4,5,6,7} ORCID- 0000-0001-8497-9903

¹ Division of Applied Biomedical Sciences and Biotechnology, School of Health Sciences, IMU University, Kuala Lumpur 57000, Malaysia

² Department of Biochemistry, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Serdang 43400, Selangor, Malaysia

³ Centre for Environmental and Population Health, Institute for Research, Development and Innovation 12 (IRDI), IMU University, 57000 Kuala Lumpur, Malaysia

⁴ British Antarctic Survey, NERC, High Cross, Madingley Road, Cambridge CB3 0ET, UK

⁵ Department of Zoology, University of Johannesburg, Auckland Park 2006, South Africa

⁶ Millennium Institute Biodiversity of Antarctic and Subantarctic Ecosystems (BASE), Las Palmeras 3425, Santiago, Chile

⁷ School of Biosciences, University of Birmingham, Edgbaston, Birmingham B15 2TT United Kingdom

* corresponding author <wongchiewyen@imu.edu.my>

Running title: Effects of heavy metals on diesel removal by algae

Abstract: Hydrocarbon spillage has long been a concern in Antarctica as it can result in detrimental effects on Antarctic biota and ecosystems. Bioremediation, using microorganisms such as microalgae, represents one of the most effective and least damaging methods developed to remove pollutants from the environment. However, the effectiveness of bioremediation in eliminating diesel can be influenced by co-contamination of the spill area by heavy metal ions, as is often the case. This study assessed the effects of zinc (Zn), lead (Pb), copper (Cu) and cadmium (Cd) on the bioremediation of diesel by a freshwater Antarctic microalga isolated from soil, *Tritostichococcus* sp. WCY_AQ5_1 (GenBank accession number: OQ225631), under laboratory conditions. Toxicity testing of heavy metals (1 to 16 ppm) on *Tritostichococcus* sp. showed that

microalgal specific growth rates and pigment ratios remained constant up till 8 ppm for all four heavy metals, an implication of toxicity at 16 ppm. In subsequent experiments, where diesel was introduced, sub-lethal Zn and Cd ion concentrations (2 to 10 ppm) did not significantly affect the biodegradation ability of *Tritostichococcus* sp. In contrast, sub-lethal Pb and Cu levels led to reduced diesel biodegradation at higher concentrations (8 to 10 ppm) by approximately 33% and 55%, respectively. Intriguingly, patterns of microalgal growth were not correlated with those of biodegradation efficiency as a prominent increase in growth of Zn-exposed cultures was observed at 8 and 10 ppm, and growth of Cu-exposed cultures peaked at 6 ppm. On the other hand, microalgal growth in Pb and Cd-exposed cultures (2 to 10 ppm) generally remained the same as control (0 ppm).

Keywords: Antarctic, Greenwich Island, polar algae, remediation, hydrocarbon pollution.

Introduction

Since the first human contact with Antarctica over two centuries ago, the region has been a magnet for marine exploitation industries, explorers, researchers and, most recently, tourists (Tin *et al.* 2014). Today, the fishing industry, shipping related with the tourism industry and the support and logistics of national research station operation and marine research in Antarctica rely almost entirely on the use of diesel hydrocarbons (Aislabie *et al.* 2004; Wong *et al.* 2021). This creates risks of accidental spillage during transfer and storage of fuel ashore, aircraft and vehicle refuelling and shipping accidents (Kerry 1993; Tin *et al.* 2010; Raymond *et al.* 2017; Wong *et al.* 2021). It has been recorded that from 1980 to 1989 alone, fuel spillages at Williams Field, an airfield in Antarctica which provides logistical support to McMurdo Station, amounted to approximately 380,000 litres (Tumeo and Larson 1994). Notably, the size of the accumulated spillage even surpassed the *Nella Dan* vessel incident which occurred in 1987 and had a spill size of 270,000 litres (Pople *et al.* 1990). Fuel leaks can also occur due to weathering and damage to containment facilities and pipelines as a result of freezing temperatures, corrosion, lack of maintenance of infrastructure at abandoned or mothballed facilities and historical waste disposal sites (Hughes and Stallwood 2005; Tin *et al.* 2010; Raymond *et al.* 2017).

Diesel is a complex mixture of approximately 10 groups of hydrocarbons (Sazhin *et al.* 2014; Wong *et al.* 2021). A large proportion typically comprises aromatic hydrocarbons, which are refractile and resistant to biodegradation due to the presence of C=C bonds (Roslee *et al.* 2021). In Antarctica, diesel persistence is further enhanced by the chronically low temperatures, which result in increased viscosity, decreased volatilisation and low water solubility (Aislabie

et al. 2006; Roslee *et al.* 2021). Inaccessible emulsions may be formed in Antarctic marine oil spill events due to strong sea surface turbulence (McGenity *et al.* 2012). The number of publications related to diesel pollution in Antarctica has been increasing with an annual growth rate of 5.84% in the past four decades, highlighting the expanding recognition of the importance of this issue (Lim *et al.* 2021)

The *Bahia Paraiso* oil spill incident in 1989, still one of the largest single oil spills to have occurred in Antarctica, led to several hundred deaths of affected penguins and shags over the subsequent three-week period primarily due to feather fouling (Kennicutt 1990; Matcott *et al.* 2019). In penguins for example, oiled feathers diminish their thermoregulatory ability and increase body drag during swimming, ultimately accelerating energy consumption in penguins while greatly limiting forage time due to increased risk of hypothermia in freezing Antarctic waters (Culik *et al.* 1991; Goldsworthy *et al.* 2000). Importantly, the refractile nature of diesel pollution as discussed earlier and its toxic properties also pose chronic threats to Antarctic biota. In particular, high and persistent aromatic content in the toxicity-determining water-accommodated fractions (WAFs) of several commonly used Antarctic fuels under polar conditions as compared with temperate conditions has been reported (Brown *et al.* 2016; Puasa *et al.* 2021). Furthermore, the high levels of polycyclic aromatic hydrocarbons (PAHs) of up to 1588 ng g⁻¹ lipid noted in fat tissues of deceased penguins is an indication of their incorporation in the Antarctic food chain (Taniguchi *et al.* 2009). While there is limited information about the association of tissue PAH concentrations with biochemical responses and genetic damage in seabirds (Albers 2006), it is well-documented that rehabilitated oiled seabirds have significantly lower post-release survival rates than their non-oiled counterparts, likely due to PAH toxicity from ingested fuel affecting their overall health (Goldsworthy *et al.* 2000). PAH toxicity have caused endocrine disruption, impaired immunity, and reproductive failure in seabirds (Franci *et al.* 2014), posing a significant threat to the restoration and maintenance of population sizes, especially among Antarctic-exclusive species.

The long-term adverse impacts of persisting residual hydrocarbon pollution highlight the necessity for remediation, which can take the form of physical, chemical and biological methods. The use of bioremediation, which eliminates environmental pollutants by utilising living organisms including microbes and plants, has gained increasing momentum in recent years as it is more efficient, cost-effective and ecologically friendly compared to other methods (Banerjee *et al.* 2016; Sharma 2020; Zakaria *et al.* 2021). Briefly, bioremediation at hydrocarbon polluted sites can take place via biosorption which involves the removal of hydrocarbons that are passively bound to the biosorbent, or more commonly, biodegradation

whereby hydrocarbons are internalised and undergo a transformation process to become harmless products (Pathak *et al.* 2018). At present, the most well-studied candidate organisms in hydrocarbon degradation are bacteria (Zakaria *et al.* 2021). However, the use of microalgae may provide several advantages including significantly reduced production of secondary waste (sludge) (Touliabah *et al.* 2022) and the potential for use in biofuel production due to the high lipid and carbohydrate content of microalgal biomass (Vo *et al.* 2018). Merit should also be given for their high carbon fixation capacity which assists in reduction of global warming (Sydney *et al.* 2014). In addition, although most microalgae cannot directly fix atmospheric nitrogen (N) to be used biologically (Coale *et al.* 2024), they are highly efficient in uptaking and assimilating inorganic N forms compared to bacteria because their phototropic nature permits them to generate energy and use water as the ultimate reductant to convert up-taken nitrates/nitrites into ammonium, the fundamental prerequisite for many biologically important molecules, in the presence of sunlight (Kumar and Bera 2020). Examples of hydrocarbon-degrading microalgae include *Chlorella vulgaris*, which is capable of removing 82–100% of oil at concentrations of 10 to 20 g/L within two weeks (Kalhor *et al.* 2017). *Nannochloropsis oculata* also showed ability to remove 66.5% of oil in oilfield produced water-containing medium within three weeks (Ammar *et al.* 2018).

Oil pollution often does not occur in isolation, and co-contamination with heavy metals is another common feature of pollution events in Antarctica (Chu *et al.* 2019). Heavy metal pollution in Antarctica is often associated with anthropogenic activities such as fuel combustion, handling and spills, and waste disposal and incineration (Goldsworthy *et al.* 2003; Chu *et al.* 2019). Natural sources of heavy metals can also contribute, from sources such as geological and glaciological weathering, wind-blown crustal dust, sea-spray, volcanism, and marine vertebrate excreta and carcasses (Chu *et al.* 2019). For instance, high levels of anthropogenically-sourced heavy metal contamination have been reported in the McMurdo Sound region of Victoria Land in Antarctica, including zinc (Zn), lead (Pb), copper (Cu) and cadmium (Cd) (Claridge *et al.* 1995) and similar findings have been reported more widely around human facilities and centres of operation (Zakaria *et al.* 2021).

Locally, Zn can be found at high levels at rubbish dump sites because of its presence in compounds commonly employed as protective coatings for fuel drums as well as in man-made accessories such as batteries, wires and nails (Claridge *et al.* 1995; Gasparon and Burgess 2000). Inappropriate disposal of Pb-containing batteries can lead to elevated concentrations of Pb in surrounding soils as their metallic layer and electrolyte solution contain PbO and Pb²⁺, respectively, while other sources include paints and cans (Claridge *et al.* 1995). High levels of

both metals have also been reported in sewage outfalls (Kennicutt *et al.* 1995). Soils near buried Cu wires and corroding Cu pipes can also contain high Cu concentrations (Claridge *et al.* 1995). Finally, transportation of Cd to Antarctica from other regions can take place through the global distribution of volatile methylated Cd generated by marine prokaryotes (Emnet 2009). Recently, analyses of Zn, Pb, Cu, and Cd in snow samples collected along transverse route from coastal research station to Antarctic ice sheet summit have shown decreasing annual deposition flux towards inland regions, suggesting that anthropogenic activities at ice-free coastal areas are a significant source of these elements (Jiang *et al.* 2018; Herath *et al.* 2024). On the other hand, elements in inland regions of Antarctica are primarily contributed by long-distance transport of anthropogenic emissions from South American continent in addition to local research stations (Herath *et al.* 2024). Importantly, higher concentrations of Pb and Cu have been observed in sea-surface microlayer than underlying water which confirm maritime Pb and Cu input from the atmosphere (Frache *et al.* 2001). It has also been suggested that maritime Zn and Cd input can occur from glacial meltwaters previously trapped in ice via atmospheric deposition (Sieber *et al.* 2019, 2020).

Although some heavy metals (*e.g.*, Zn and Cu) are essential to microalgae, exposure to heavy metals at concentrations above their maximum tolerance level will incur negative impacts (Monteiro *et al.* 2012). For instance, Zn concentrations below 15 ppm improved growth of *Dunaliella tertiolecta* while at 25 ppm the microalga greatly suffered from toxic effects (El-Agawany and Kaamouh 2023). Generally, excessive Zn can induce toxicity in microalgae by disrupting membrane integrity, altering intracellular sodium and potassium concentrations, and inhibiting photosynthesis (Wong and Chau 1990). Pb toxicity is often associated with diminished photosynthetic ability, elevated oxidative stress and defects in cell division in microalgae (Küpper 2017). Microalgae can also experience reduced ability to photosynthesise and respire, at the same time develop organelles of abnormal size and morphology, when exposed to high Cu concentrations, thus reducing growth rates (Rocha *et al.* 2021). Exposure to elevated Cd concentration can lead to dilation of membrane-containing organelles, granule accumulation and the substitution of essential metal components in enzymes in microalgae (Vymazal 1987).

Heavy metal exposure can also alter the way that cells and hydrocarbon fractions interact, as shown by PAH biodegradation rates best fitting first-order kinetics when heavy metals are present in a study performed with soil microcosms (Baltrons *et al.* 2018). Previously, it was also reported that biodegradation of low molecular weight PAHs by green microalga, *Selenastrum capricornutum*, were significantly enhanced under exposure to a mixture of heavy

metals (Ke *et al.* 2010). It is therefore crucial to understand the role that heavy metals play during hydrocarbon biodegradation (Lim *et al.* 2021). As *Tritostichococcus* sp. WCY_AQ5_1 is the first Antarctic green microalga confirmed to possess diesel degradation ability (Lim *et al.* 2023), the primary aim of the current study was to assess the impact of the heavy metal ions, Zn, Pb, Cu and Cd, on this Antarctic microalga and how these co-contaminants affect its ability to bioremediate diesel, based on changes in microalgal growth, pigment content, oxidative stress responses and diesel biodegradation efficiency.

Material and methods

Study alga and culture conditions. — The freshwater green unicellular microalgal strain, *Tritostichococcus* sp. WCY_AQ5_1 (GenBank accession number: OQ225631) used in this study was originally isolated from soil samples obtained from Greenwich Island (South Shetland Islands; 62.4692°S, 59.7963°W) (Lim *et al.* 2023). The alga was grown in optimised Bold's Basal Medium (BBM) and maintained in an incubator at 10°C (Lim *et al.* 2023) under illumination with cool white fluorescent lamps (Philips, TLD 18W/54-765 providing 42 $\mu\text{mol m}^{-2} \text{s}^{-2}$ PAR) on a 12h:12h light:dark cycle.

Heavy metals and diesel source. — Atomic absorption spectroscopy grade heavy metal stock solutions containing 1000 ppm of one of the following heavy metals: zinc (Zn), lead (Pb), copper (Cu), or cadmium (Cd), were obtained from Sigma-Aldrich Chemicals, USA. Dilution was used to obtain the required heavy metal test concentrations using optimised BBM. Petronas diesel (Diesel Euro B10) was purchased from a local fuel station in Kuala Lumpur. The diesel was filter-sterilised using an 0.22 μm nylon membrane filter and stored in an amber Scott bottle at room temperature.

Toxicity testing of heavy metals. — In initial trials, the microalga was exposed to a broad range of heavy metal concentrations to establish a suitable test concentration range. An exponential phase inoculum of *Tritostichococcus* sp. was standardised at OD₆₂₀ (absorbance value) to 0.5. Fifty millilitre cultures were grown in 100 mL flasks consisting of 20% microalgae and one of each of the heavy metal standard solutions (Zn, Pb, Cu, Cd), with concentrations varying across the range 0, 1, 2, 4, 8, 16 ppm. Incubation took place under controlled conditions as described in Section 2.1 for 7 d. During this period, microalgal growth was assessed on days 0, 1, 3, 5 and 7 through determination of chlorophyll *a* (chl-*a*) content.

Determination of photosynthetic pigment contents. — As a measure of algal growth, the concentrations of chl-*a*, chlorophyll *b* (chl-*b*) and carotenoids (car) in the microalgal

cultures on day 7 were determined by spectrophotometry after methanol extraction. Five mL of resuspended culture from each flask were harvested and centrifuged at 4,000 rpm at 4°C for 20 min. The cell pellet was resuspended in 2 mL analytical grade 100% methanol (Sigma-Aldrich, USA) and sonicated in iced water for 1 h. The resulting samples were stored in the dark for 24 h at 4°C, before being centrifuged at 4,000 rpm and 4°C for 25 min followed by OD measurement of the supernatants at 665.2, 652.4 and 470 nm. Concentrations of chl-*a*, chl-*b*, and car were calculated using the following equations (Lichtenthaler and Buschmann 2001):

$$\begin{aligned} \text{Chl} - a \text{ (mg/L)} &= 16.72A_{665.2} - 9.16A_{652.4} \\ \text{Chl} - b \text{ (mg/L)} &= 34.09A_{652.4} - 15.28A_{665.2} \\ \text{Car (mg/L)} &= \frac{1000A_{470} - 1.63 \times (\text{Chl} - a) - 104.96 \times (\text{Chl} - b)}{221} \end{aligned}$$

Determination of specific growth rate. — Using the values of chl-*a* obtained, the microalgal specific growth rates (μ) expressed in unit reciprocal days (d^{-1}) under each heavy metal treatment concentration were determined using the following formula (Krzemińska *et al.* 2014):

$$\mu \text{ (d}^{-1}\text{)} = \frac{\ln N_t - \ln N_0}{t - t_0}$$

where N_t and N_0 represent chl-*a* concentrations at time t and t_0 respectively.

Assessment of the influence of heavy metals on diesel hydrocarbon bioremediation.

— Fifty millilitres of microalgal culture were grown in 100 mL flasks containing 20% microalgal inoculum ($OD_{620} = 0.5$), 1% v/v diesel, and a single heavy metal standard solution (Zn, Pb, Cu, Cd) at concentrations 0, 2, 4, 6, 8 or 10 ppm. An abiotic control test was run using optimised BBM and 1% v/v diesel. Aeration with syringe-filtered ambient air was implemented for all cultures and incubation took place under controlled conditions as described in Section 2.1 for 7 d. Biodegradation efficiency was measured at the end of the experiment, and a sub-sample of culture was harvested for the determination of growth by assessment of photosynthetic pigment content (Section 2.3.1), and measurement of oxidative stress responses.

Assessment of microalgal diesel biodegradation efficiency. — To measure biodegradation efficiency, the experimental microalgal cultures were subjected to *n*-hexane extraction (1:1 medium to solvent ratio). The mixtures were thoroughly swirled and allowed to settle for 10 min until two separated layers were formed (culture layer and *n*-hexane layer). The *n*-hexane layer containing residual diesel was recovered onto a pre-weighed Petri dish followed by 24 h drying under a fume hood at room temperature to ensure complete vaporisation of the extraction solvent. The residual diesel mass was obtained via subtracting

the original mass of the Petri dish from its final mass and the % biodegradation efficiency (BE) of the microalga was calculated as follows (Chen *et al.* 2013; McFarlin *et al.* 2018):

$$BE\% = \frac{(\text{Residual diesel in abiotic control (g)} - \text{Residual diesel in experimental cultures(g)})}{\text{Diesel introduced (g)}} \times 100$$

Determination of reactive oxygen species levels. — Reactive oxygen species (ROS) analysis was carried out using the cell permeable indicator 2, 7-dichlorodihydrofluorescein diacetate (H2DCFDA) (Invitrogen™, Thermo Fisher Scientific, USA). Two hundred microlitres of experimental culture were added to 4 µL of 20 mM H2DCFDA in a 96-well Nucleon black plate (ThermoFisher Scientific, USA) in triplicates and resuspended thoroughly. The plate was left in the dark for 15 min at room temperature. Then, fluorescence was measured at 495 nm excitation and 530 nm emission wavelengths using a microplate reader (Tecan Infinite F200).

Determination of lipid peroxidation. — Lipid peroxidation (LP) analysis used the cell permeable indicator C11-BODIPY^{581/591} (Life Technologies®, USA). Two hundred microlitres of experimental culture were added to 5 µL of 5 mM C11-BODIPY^{581/591} in a 96-well Nucleon black plate in triplicates and resuspended thoroughly. The plate was left in the dark for 30 min at room temperature. Fluorescence was then measured at 485 nm excitation and 530 nm emission wavelengths using a microplate reader.

Statistical analyses. — All statistical analyses were performed using GraphPad Prism 9. Means and standard deviations were calculated and compared using one-way analysis of variance (ANOVA) and, where significant, pair-wise Tukey *post-hoc* tests were then carried out. Differences were considered significantly different at $p < 0.05$. Where non-parametric tests were necessary, Kruskal-Wallis test was applied.

Results

Toxicity testing of heavy metals on *Tritostichococcus* sp. WCY_AQ_1

Growth response. — As shown in Fig. 1, results suggest that exposure of microalgal cultures to heavy metal concentrations from 1 to 8 ppm, regardless of the type of heavy metal, does not negatively impact cell growth because their growth trends largely overlapped with the trend exhibited by their respective control groups. In line with this, although one-way ANOVA revealed significant differences in the μ over 7 d among cultures subjected to different concentrations of Zn ($F = 22.94$, $p < 0.0001$), Pb ($F = 6.017$, $p = 0.0052$), or Cu ($F = 222.5$, $p < 0.0001$), subsequent Tukey's *post-hoc* tests revealed that the microalgal cultures only showed a significant drop in their μ when exposed to these heavy metals at 16 ppm (Table 1), relative to

μ obtained at other test concentrations, hence inferring toxicity at 16 ppm. In contrast, comparatively good tolerance of exposure to Cd was apparent up to 16 ppm, with μ values not differing significantly across the concentrations tested.

Photosynthetic pigment content. — No significant changes in chl-*a*, chl-*b* or car contents were observed under exposure to 0–8 ppm concentrations all four heavy metals (Fig. 2). The pigment ratio, chl-*b*/chl-*a*, remained relatively constant across all concentrations tested, suggesting that any changes in chl-*b* and chl-*a* content were of similar magnitude. However, one-way ANOVA identified significant differences in the car/chl-*a* ration of microalgal cells when subjected to different heavy metal test concentrations (Zn: $F = 20.23$, $p < 0.0001$; Pb: $F = 17.49$, $p < 0.0001$; Cu: $F = 46.38$, $p < 0.0001$; Cd: $F = 22.61$, $p < 0.0001$). Subsequent Tukey's *post-hoc* tests confirmed that only at 16 ppm were the car/chl-*a* ration of microalgal cells significantly lower than cultures tested with lower heavy metal concentrations (0–8 ppm) and this observation was consistent for all four heavy metals.

Effect of heavy metals on diesel bioremediation by *Tritostichococcus* sp. WCY_AQ5_1

Growth response and biodegradation efficiency. — In the presence of diesel, the microalgal cultures exposed to Zn, Pb and Cd showed overall lower chl-*a* concentration ranges on day 7 (Zn: 0.398–0.879 mg/L, Pb: 0.364–0.519 mg/L, Cd: 0.169–0.221 mg/L) as compared to those of cultures exposed to heavy metals alone (Zn: 1.128–1.268 mg/L, Pb: 0.908–1.087 mg/L, Cd: 0.696–0.777 mg/L) (Fig. 3). A similar pattern was seen in the combined diesel and Cu treatment, except for cultures exposed to Cu at 6 ppm where a peak in growth (Fig. 4C) was seen which surpassed those achieved with Cu treatment alone (Fig. 2C).

One-way ANOVA comparing the effects of different heavy metal concentrations on microalgal growth revealed significant differences in growth response for diesel-exposed cultures under Zn ($F = 49.53$, $p < 0.0001$) and Cu ($F = 39.71$, $p < 0.0001$) treatment. Tukey's *post-hoc* test results further revealed that Zn significantly enhanced microalgal growth at 8 and 10 ppm compared to other test concentrations (Fig. 4A). In cultures exposed to Cu, microalgal growth was significantly greater at 6 ppm than at the other concentrations tested (Fig. 4C). Cu concentrations above 6 ppm resulted in reduced growth. Pb and Cd concentration did not have a significant influence on microalgal growth across all concentrations tested. Nevertheless, as shown in Fig. 4D, increasing Cd concentration was associated with reduced microalgal growth.

One-way ANOVA also identified significant effects of different heavy metal concentrations on biodegradation efficiency (BE, %) (Pb: $F = 17.91$, $p = 0.0004$, Cu: $F = 16.87$, $p = 0.0005$). *Post-hoc* pairwise tests confirmed significantly lower BE in cultures exposed to 8

and 10 ppm Pb (Fig. 5B) and Cu (Fig. 5C) as compared to groups exposed to lower concentrations. However, BE of cultures exposed to different concentrations of Zn and Cd were not significantly different although, again, a slight reduction in BE with increasing Cd concentration can be observed in Fig. 4D.

Photosynthetic pigment content. — One-way ANOVA confirmed significant differences in all three pigment contents in diesel-exposed microalgal cells when subjected to different concentrations of Zn (Fig. 5A) (chl-*a*: $F = 49.53$, $p < 0.0001$; chl-*b*: $F = 18.88$, $p < 0.0001$; car: $F = 9.308$, $p = 0.0011$) and Cu (Fig. 4c) (chl-*a*: $F = 39.71$, $p < 0.0001$; chl-*b*: $F = 26.52$, $p < 0.0001$; car: $F = 78.51$, $p < 0.0001$), but no significant differences for Pb (Fig. 5B) and Cd-treated (Fig. 5D) cultures. The changes in individual photosynthetic pigment content (Fig. 5) were consistent with the measure of cell growth (Fig. 4). In addition, one-way ANOVA of pigment ratios showed significant changes in the chl-*b*/chl-*a* ($F = 10.65$, $p = 0.0009$) ratio of Zn-treated cultures, and in both chl-*b*/chl-*a* ($F = 5.179$, $p = 0.0163$) and car/chl-*a* ($F = 29.77$, $p < 0.0001$) ratios for Cu-treated cultures in the presence of diesel, with a peak at 2 ppm (Fig. 5A) and 0 ppm (Fig. 5C), respectively; minimal changes in chl-*b*/chl-*a* and car/chl-*a* ratios were observed when treatment concentrations of Zn and Cu were further increased.

Oxidative stress response. — Figure 6 illustrates the ROS and LP levels in microalgal cells after exposure to diesel in combination with various concentrations of different heavy metals. One-way ANOVA revealed significant differences in ROS for diesel-exposed cultures subjected to different concentrations of Zn (Fig. 6A) ($F = 23.48$, $p < 0.0001$), Cu (Fig. 6C) ($F = 204.8$, $p < 0.0001$) and Cd (Fig. 6D) ($F = 68.22$, $p < 0.0001$), while significant differences in LP were only observed for those exposed to Zn (Fig. 6E) ($F = 78.65$, $p < 0.0001$) and Cu (Fig. 6G) ($F = 695.6$, $p < 0.0001$). In Cu-treated cultures, ROS and LP levels decreased as Cu concentration increased (Figs. 6C, G). In contrast, increasing ROS levels (Fig. 6A) were found in combination with decreasing LP levels (Fig. 6E) in Zn-exposed samples. Cells subjected to Cd treatment showed increased ROS levels as the concentration of Cd increased (Fig. 6D) but with no significant changes in LP levels (Fig. 6H). Pb treatment resulted in ROS and LP levels that did not differ significantly from those observed in the control (0 ppm) (Figs. 6B, F).

Discussion

Zn and Cu are essential micronutrients for algal growth as they are components of a diverse group of enzymes required for algal metabolism and photosynthesis-associated electron transport reactions (Coleman 1998; Raven *et al.* 1999). Conversely, Pb and Cd are non-essential

metals and are considered as toxic. Notably, pigment concentrations and ratios of *Tritostichococcus* sp. subjected to 1–8 ppm of Zn, Pb, Cu or Cd showed no significant differences from the control group and no metal-specific effects (Fig. 2). Nonetheless, co-treatment of microalgal cultures with sublethal heavy metal concentrations (1–10 ppm) and diesel generally resulted in lower growth response as compared to that under heavy metal treatment alone (Fig. 3) as well as noticeable changes in microalgal growth patterns (Fig. 4).

The drastically suppressed microalgal growth (Fig. 3) could indicate diesel toxicity, as a negative correlation between diesel exposure and chl-*a* content has previously been reported in *Pseudokirchneriella subcapitata* and *Chlorella* sp. MM3 (Ramadass *et al.* 2017). The study also noted greater toxicity induced by the diesel WAF, *i.e.*, the water-soluble proportion of diesel that were made bioavailable to organisms, as compared to diesel itself. Other mechanisms involved could include reduced light penetration as a result of slick formation, impaired photosynthetic ability due to membrane distortion, and extracellular leaching of chlorophyll (O'Brien and Dixon 1976; González *et al.* 2009).

The presence of some heavy metals may alleviate diesel-induced toxicity experienced by microalgal cells. Our data suggest Zn may play such a role, as significant growth of microalgal cells was observed when exposed to 8 or 10 ppm of Zn in combination with diesel (Fig. 4A). Consistent with this, a marked decrease in LP was observed in microalgal cells at higher Zn concentrations (Fig. 6E) despite the concomitant increment in ROS levels (Fig. 6A). Many polar algae, such as *Ulothrix zonata* and Antarctic strains of *Stichococcus* sp., have been reported to possess high proportions of unsaturated fatty acids in their membranes as compared to temperate algae, highlighting their importance in maintenance membrane fluidity at low temperatures (Osipova *et al.* 2009; Chen *et al.* 2012). Given that unsaturated fatty acid chains are the primary targets of ROS attack (Rezayian *et al.* 2019), the antioxidative defence system of polar microalgae may be robust, as low temperatures have been observed to induce expression of glutathione *S*-transferase, an efficient antioxidative enzyme, in *Chlamydomonas* sp. and other Antarctic microbes (Kan *et al.* 2006; Hou *et al.* 2019). Zn treatment has also been observed to increase synthesis of superoxide dismutase (Coleman 1998) and upregulate ROS scavenging enzyme activities of peroxidases in *Chlorella sorokiniana* and *Scenedesmus acuminatus* (Hamed *et al.* 2017). Notably, it has been reported that, unlike temperate microalgae and non-photosynthetic polar microorganisms, polar microalgae have a higher demand for Zn as they hold greater diversity of Zn-binding proteins involved in regulating photosynthesis-related processes and other primary metabolism (Ye *et al.* 2022). In our study, it is possible that much of the Zn supplied was utilised in processes that combat ROS instead of diesel catabolism,

thereby causing the lack of change in BE despite increased microalgal growth at 8 and 10 ppm Zn (Fig. 4A).

An increasing ability of microalgal cells to overcome diesel toxicity in the presence of heavy metals could also be observed in Cu-treated cultures but at lower concentrations (0–4 ppm) (Fig. 4C). Combined diesel and Cu treatment at a higher concentration of 6 ppm resulted in higher chl-*a* concentration (1.353 mg/L) in microalgal cells than those subjected to Cu only (0.843–0.883 mg/L). This phenomenon could be due to the synergistic effects of diesel and Cu in a co-contaminated environment which can enhance the secretion of extracellular polymeric substances (EPS) (Ali *et al.* 2022). EPS functions as a protective barrier that prevents direct exposure of the microalga to diesel by neutralising cell surface charges, thereby enhancing the hydrophobicity of microalgal cells towards diesel whilst evading its toxic effects (Wang *et al.* 2008; Ali *et al.* 2022). However, the interactive chemistry between microalgae and co-contaminants has received little study. Further increase in Cu concentrations up to 8 ppm and 10 ppm in diesel-containing culture resulted in decreased microalgal growth (Fig. 4C). The toxic effects of Cu may arise at high concentrations through oxidative stress, as oxidation of Cu produces ROS which can in turn lead to degradation of photosynthetic pigments, oxidation of proteins and membrane lipids and DNA damage (Sujetovienė 2014; Kaamouh *et al.* 2022). However, such a mechanism is unlikely in the current study given the reduced ROS and LP levels observed at higher Cu concentrations (Figs. 6C, G). Instead, high Cu levels may have resulted in reduced permeability of the microalgal cell membrane and impeded the binding of other essential microelements such as manganese that also serve as cofactors in photosynthesis (Sunda and Huntsman 1983).

In contrast with Zn and Cu, addition of the non-essential heavy metals, Pb and Cd, to diesel-containing culture resulted in insignificant impact on microalgal growth at the concentrations tested (Fig. 3). Oxidative stress response tests were consistent with this observation, as ROS and LP levels were not significantly different to those observed in the control cultures (Figs. 6B, F), implying absence of impact of Pb treatment up to 10 ppm. It is notable, however, that BE dropped significantly under exposure to Pb at concentrations of 8 and 10 ppm (Fig. 4B). This could be a result of the powerful enzyme-binding ability of Pb causing enzyme activity inhibition (Souza *et al.* 2012). Supporting this suggestion, dehydrogenase activity in soil has been shown to be inhibited in areas with high Pb contamination (Łukowski and Dec 2021). Dehydrogenase is one of the main classes of microbial enzyme involved in oxidative activities, including the catabolism of diesel hydrocarbons into organic carbon-containing metabolites (Errington *et al.* 2018; Łukowski and

Dec 2021). Other enzymes which may be affected include dioxygenases, an important enzyme group that plays a role in bio-transforming benzo[a]pyrene hydrocarbon and has been recently reported to be present in the hydrocarbon-degrading alga, *Selenastrum capricornutum* (García de Llasera *et al.* 2022). Other than the protein level, Pb could also interfere with the transcription of hydrocarbon-degrading enzymes. In bacteria, for example, cytochrome P108J1 has previously been reported to play an important role in the initial breakdown of PAHs (Luo *et al.* 2016). Since Pb and some other heavy metals have also been shown to alter the expression of cytochrome P450 genes, and an increasing library of cytochrome p450 genes have been identified in microalgae, Pb may play a role in downregulating the expression of these genes resulting in degradation of hydrocarbons becoming less effective (Korashy and El-Kadi 2005; Zheng *et al.* 2022). However, functional studies on these microalgal enzymes remain limited (Zheng *et al.* 2022).

In comparison with the other heavy metals tested, the impact of Cd addition to diesel-containing cultures were minimal in terms of both microalgal growth and BE (Fig. 4D). Cd is generally regarded as toxic towards microalgae as it leads to mitochondrial damage, disrupts cell motility, affects cellular components involved in photosynthesis, and inhibits protein synthesis and growth of microalgae (Trevors *et al.* 1986). Nevertheless, the current study demonstrated relatively strong tolerance of *Tritostichococcus* sp. towards Cd, with LP levels that differed minimally from the control treatment across all tested concentrations in cultures containing diesel (Fig. 6H) and insignificant impact on μ resulting from Cd addition up to 16 ppm (Fig. 1D). As Cd has been reported to be one of the most effective inducers of phytochelatin in microalgae (Ahner and Morel 1995), increased production of phytochelatin may be a possible mechanism that contributes to such strong tolerance as it enhances ROS scavenging, prevents heavy metal build-up in the cytoplasm and the non-specific binding of heavy metals to important biomolecules by sequestering the peptide-bound heavy metals into the microalgal vacuole (Scarano and Morelli 2002; Tsuji *et al.* 2002). In addition, good metal-phytochelatin binding properties of Cd (>80%) in *Phaeodactylum tricornerutum* compared to other non-essential heavy metals such as Pb (40%) have previously been reported (Scarano and Morelli 2002). Nonetheless, there was a noticeable but non-significant gradual decrease in both chl-*a* and BE with increasing Cd concentration (Fig. 4D). Similar findings have been reported in *Rhodococcus* sp., whereby inhibitory effects of Cd were reflected in the growth of bacteria and their oil-degrading ability (Ibrahim *et al.* 2020). This phenomenon could be due to the slight downregulation of antioxidative enzyme cofactors such as glutathione, which have a role in phytochelatin formation (Tsuji *et al.* 2003; Zhao *et al.* 2019).

In comparison to other studies which assessed the impact of various heavy metals on hydrocarbon degradation ability of Antarctic microbes, it was previously reported that 1 ppm of Zn and Cd were sufficient to reduce the degradation of waste canola oil by a bacterial community isolated near an Antarctic research station by approximately 40% and 25%, respectively (Zahri *et al.* 2020). In another similar study which used the same bacterial community, 1 ppm of Cu also slightly hindered degradation of waste canola oil compared to control whereas the inhibitory effects of Pb were observed at 10 ppm (Zahri *et al.* 2021). *Rhodococcus* sp. (AQ5-07) which waste canola oil degradation ability was unaffected by 1 ppm of Zn, Pb, and Cu, was significantly influenced at 0.4 ppm of Cd concentration with an approximate 20% drop in oil removed (Ibrahim *et al.* 2020). For a fair comparison considering the type of hydrocarbon being degraded, diesel degradation ability of an undefined Antarctic bacterium strain AQ5-AO1 was significantly impaired with exposure to Zn and Cd at 1 ppm (Zakaria *et al.* 2020). Degradation of phenanthrene, a PAH, was almost completely inhibited at 5 ppm of Cd in another Antarctic bacterium, *Sphingobium xenophagum* (Gran-Scheuch *et al.* 2017). In this respect, *Tritostichococcus* sp. has outperformed many reported Antarctic bacteria in maintaining control (0 ppm)-comparable BE up to 10 ppm for Zn and Cd-exposed cultures, and 6 ppm for Cu-exposed cultures whereas tolerance to Pb was similar for the bacterial community reported by Zahri *et al.* (2021). The same studies also observed that bacterial growth was generally inhibited at around 1–5 ppm of Zn, Pb, Cu, and Cd (Gran-Scheuch *et al.* 2017; Ibrahim *et al.* 2020; Kai *et al.* 2020; Zahri *et al.* 2020; Zakaria *et al.* 2020). This suggests that hydrocarbon-degrading Antarctic bacteria are more sensitive to heavy metal interference than *Tritostichococcus* sp., where growth inhibition only became evident above 6 ppm of Cu (Fig. 4C) whereas impacts of other heavy metals were either minimal (Figs. 4B, D) or even positive on growth (Fig. 3A).

Content of heavy metals in soils of Antarctic anthropogenically-affected sites can differ drastically with range 11.3–116.2 ppm for Zn, 0.05–68.7 ppm for Pb, 2.56–145.2 ppm for Cu, and 0.005–0.308 ppm for Cd reported in Larsemann Hills and Mirny station (Alekseev and Abakumov 2021). In another study, samples from topsoil of Mirror Peninsula that is exposed to a wide range of weathering processes, anthropogenic impacts, and biological activity ranged 1.05–15.3 ppm for Zn, 6.68–98.9 ppm for Pb, and 0.071–24.8 ppm for Cu (Xu *et al.* 2020). Sediments in vicinity of a research station from Hangar Cove, South Cove, and Lagoon Island found range 29.6–54.3 ppm for Zn, 4.8–5 ppm for Pb, 19.6–44.3 ppm for Cu, and 0.2–0.5 ppm for Cd (Webb *et al.* 2020). Meanwhile, positive correlation was found between organic matter and levels of Zn and Cd in soil samples collected from Admiralty Bay, indicating that selective

enrichment of these heavy metals is also influenced by the presence of marine vertebrates and their feces (Souza-Kasprzyk *et al.* 2022). Given the substantial difference in heavy metal concentrations within each site and from site-to-site, the higher the microbial tolerance towards heavy metals, the higher the chances for the microbe to thrive and subsequently carry out bioremediation activities in various environmental conditions. This preliminary study has shown outstanding ability of *Tritostichococcus* sp. to withstand Zn, Pb, Cu, and Cd at concentrations that are several folds of other reported Antarctic hydrocarbon-degrading microbes to remove diesel, making them promising bioremediation candidates for the co-contaminated Antarctic land.

Conclusions

The current study confirmed that different heavy metals have different impacts on *Tritostichococcus* sp. WCY_AQ5_1, reflected in changes in microalgal growth and diesel degradation ability. Cultures supplemented with the essential heavy metals, Zn and Cu, generally demonstrated improved microalgal growth in diesel-containing medium and these metals likely played a role in limiting diesel toxicity. Conversely, the influence of Pb on microalgal growth was insignificant although BE was reduced at higher concentrations. The microalga showed strong tolerance of exposure to Cd even at the highest concentrations tested here compared to the other tested heavy metals, which caused only a slight decline in microalgal growth and BE. Overall, the microalga demonstrated relatively robust survival resilience and retention of diesel biodegradation ability compared to other existing Antarctic hydrocarbon-degrading microbes in an environment where considerable amount of heavy metals have been added. Moreover, these promising results were obtained under controlled laboratory conditions with temperatures slightly higher than those typically found *in situ*. This suggests that *Tritostichococcus* sp. might show optimal bioremediation activity during the Antarctic summer or potentially for extended periods annually due to increasing global warming.

Acknowledgements. — The authors would like to acknowledge the internal grant (Grant No. MBT I-2022 (04)) from the IMU University and national grant (Grant No. IMU R 271/2021) from the Yayasan Penyelidikan Antartika Sultan Mizan (YPASM) Research Grant 2020 in supporting the completion of this research project. Peter Convey is supported by NERC core funding to the British Antarctic Survey's 'Biodiversity, Evolution and Adaptation' Team. Special thanks to the reviewers for their instrumental feedback which has greatly improved the quality of this manuscript.

References

- Ahner B.A. and Morel F.M.M. 1995. Phytochelatin production in marine algae. 2. Induction by various metals. *Limnology and Oceanography* 40: 658–665. doi: 10.4319/lo.1995.40.4.0658
- Aislabie J.M., Balks M.R., Foght J.M. and Waterhouse E.J. 2004. Hydrocarbon spills on Antarctic soils: Effects and management. *Environmental Science and Technology* 38: 1265–1274. doi: 10.1021/es0305149
- Aislabie J., Saul D.J. and Foght J.M. 2006. Bioremediation of hydrocarbon-contaminated polar soils. *Extremophiles* 10: 171–179. doi: 10.1007/s00792-005-0498-4
- Albers P.H. 2006. Birds and polycyclic aromatic hydrocarbons. *Avian and Poultry Biology Reviews* 17: 125–140. doi: 10.3184/147020606783438740
- Alekseev I. and Abakumov E. 2021. Content of trace elements in soils of Eastern Antarctica: Variability across landscapes. *Archives of Environmental Contamination and Toxicology* 80: 368–388. doi: 10.1007/s00244-021-00808-4
- Ali M., Song X., Ding D., Wang Q., Zhang Z. and Tang Z. 2022. Bioremediation of PAHs and heavy metals co-contaminated soils: Challenges and enhancement strategies. *Environmental Pollution* 295: 118686. doi: 10.1016/J.ENVPOL.2021.118686
- Ammar S.H., Khadim H.J. and Mohamed A.I. 2018. Cultivation of *Nannochloropsis oculata* and *Isochrysis galbana* microalgae in produced water for bioremediation and biomass production. *Environmental Technology and Innovation* 10: 132–142. doi: 10.1016/J.ETI.2018.02.002
- Baltrons O., López-Mesas M., Vilaseca M., Gutiérrez-Bouzán C., Le Derf F., Portet-Koltalo F. and Palet C. 2018. Influence of a mixture of metals on PAHs biodegradation processes in soils. *Science of the Total Environment* 628–629: 150–158. doi: 10.1016/j.scitotenv.2018.02.013
- Banerjee A., Dutta S., Mondal S. and Roy A. 2016. Bioremediation of hydrocarbon - A review. *International Journal of Advanced Research* 4: 1303–1313. doi: 10.21474/IJAR01/734
- Brown K.E., King C.K., Kotzakoulakis K., George S.C. and Harrison P.L. 2016. Assessing fuel spill risks in polar waters: Temporal dynamics and behaviour of hydrocarbons from Antarctic diesel, marine gas oil and residual fuel oil. *Marine Pollution Bulletin* 110: 343–353. doi: 10.1016/j.marpolbul.2016.06.042
- Chen Q., Bao M., Fan X., Liang S. and Sun P. 2013. Rhamnolipids enhance marine oil spill bioremediation in laboratory system. *Marine Pollution Bulletin* 71: 269–275. doi: 10.1016/J.MARPOLBUL.2013.01.037
- Chen Z., He C. and Hu H. 2012. Temperature responses of growth, photosynthesis, fatty acid and nitrate reductase in Antarctic and temperate *Stichococcus*. *Extremophiles* 16: 127–133. doi: 10.1007/s00792-011-0412-1
- Chu W.L., Dang N.L., Kok Y.Y., Yap I.K.S., Phang S.M. and Convey P. 2019. Heavy metal pollution in Antarctica and its potential impacts on algae. *Polar Science* 20: 75–83.

- doi: 10.1016/J.POLAR.2018.10.004
- Claridge G.G.C., Campbell I.B., Powell H.K.J., Amin Z.H. and Balks M.R. 1995. Heavy metal contamination in some soils of the McMurdo Sound region, Antarctica. *Antarctic Science* 7: 9–14. doi: 10.1017/S0954102095000034
- Coale T.H., Loconte V., Turk-Kubo K.A., Vanslebrouck B., Mak W.K.E., Cheung S., Ekman A., Chen J.H., Hagino K., Takano Y., Nishimura T., Adachi M., Le Gros M., Larabell C. and Zehr J.P. 2024. Nitrogen-fixing organelle in a marine alga. *Science* 384: 217–222. doi: 10.1126/SCIENCE.ADK1075/SUPPL_FILE/SCIENCE.ADK1075_MOVIES_S1_TO_S3.ZIP
- Coleman J.E. 1998. Zinc enzymes. *Current Opinion in Chemical Biology* 2: 222–234. doi: 10.1016/S1367-5931(98)80064-1
- Culik B.M., Wilson R.P., Woakes A.T. and Sanudo F.W. 1991. Oil pollution of Antarctic penguins: Effects on energy metabolism and physiology. *Marine Pollution Bulletin* 22: 388–391. doi: 10.1016/0025-326X(91)90341-O
- El-Agawany N.I. and Kaamouh M.I.A. 2023. Role of zinc as an essential microelement for algal growth and concerns about its potential environmental risks. *Environmental Science and Pollution Research* 30: 71900–71911. doi: 10.1007/s11356-022-20536-z
- Emnet P. 2009. *Heavy metals: A heavy burden on the icy continent*. Thesis in Postgraduate Certificate in Antarctic Studies. University of Canterbury.
- Errington I., King C.K., Wilkins D., Spedding T. and Hose G.C. 2018. Ecosystem effects and the management of petroleum-contaminated soils on subantarctic islands. *Chemosphere* 194: 200–210. doi: 10.1016/J.CHEMOSPHERE.2017.11.157
- Frache R., Abelmoschi M.L., Grotti M., Ianni C., Magi E., Soggia F., Capodaglio G., Turetta C. and Barbante C. 2001. Effects of ice melting on Cu, Cd and Pb profiles in Ross Sea waters (Antarctica). *International Journal of Environmental Analytical Chemistry* 79: 301–313. doi: 10.1080/03067310108044391
- Franci C.D., Guillemette M., Pelletier É., Chastel O., Bonnefoi S. and Verreault J. 2014. Endocrine status of a migratory bird potentially exposed to the Deepwater Horizon oil spill: A case study of northern gannets breeding on Bonaventure Island, Eastern Canada. *Science of the Total Environment* 473–474: 110–116. doi: 10.1016/j.scitotenv.2013.12.006
- García de Llasera M.P., Fuentes Pérez A.C., Peralta Marín G. and Beltrán Calva E.G. 2022. First evidence of extracellular enzymatic degradation of benzo(a)pyrene by the phytoplankton species *Selenastrum capricornutum* and the influence of temperature. *Environmental Advances* 8: 100246. doi: 10.1016/j.envadv.2022.100246
- Gasparon M. and Burgess J.S. 2000. Human impacts in Antarctica: Trace-element geochemistry of freshwater lakes in the Larsemann Hills, East Antarctica. *Environmental Geology* 39: 963–976. doi: 10.1007/s002549900010

- Goldsworthy P.M., Canning E.A. and Riddle M.J. 2003. Soil and water contamination in the Larsemann Hills, East Antarctica. *Polar Record* 39: 319–337.
doi: 10.1017/s003224740300305x
- Goldsworthy S.D., Giese M., Gales R.P., Brothers N. and Hamill J. 2000. Effects of the *Iron Baron* oil spill on little penguins (*Eudyptula minor*). II. Post-release survival of rehabilitated oiled birds. *Wildlife Research* 27: 573–582. doi: 10.1071/WR99076
- González J., Figueiras F.G., Aranguren-Gassis M., Crespo B.G., Fernández E., Morán X.A.G. and Nieto-Cid M. 2009. Effect of a simulated oil spill on natural assemblages of marine phytoplankton enclosed in microcosms. *Estuarine, Coastal and Shelf Science* 83: 265–276.
doi: 10.1016/j.ecss.2009.04.001
- Gran-Scheuch A., Fuentes E., Bravo D.M., Jiménez J.C. and Pérez-Donoso J.M. 2017. Isolation and characterization of phenanthrene degrading bacteria from diesel fuel-contaminated Antarctic soils. *Frontiers in Microbiology* 8: 1634. doi: 10.3389/fmicb.2017.01634
- Hamed S.M., Zinta G., Klöck G., Asard H., Selim S. and AbdElgawad H. 2017. Zinc-induced differential oxidative stress and antioxidant responses in *Chlorella sorokiniana* and *Scenedesmus acuminatus*. *Ecotoxicology and Environmental Safety* 140: 256–263.
doi: 10.1016/j.ecoenv.2017.02.055
- Herath I.K., Ma H., Bargagli R., Hu Y., Hirabayashi M., Motoyama H. and Shi G. 2024. Sources and distribution of trace elements in surface snow from coastal Zhongshan Station to Dome A (East Antarctica). *Atmospheric Environment* 331: 120583.
doi: 10.1016/J.ATMOSENV.2024.120583
- Hou Y., Qiao C., Wang Y., Wang Y., Ren X., Wei Q. and Wang Q. 2019. Cold-adapted glutathione S-transferases from Antarctic psychrophilic bacterium *Halomonas* sp. ANT108: Heterologous expression, characterization, and oxidative resistance. *Marine Drugs* 17: 147.
doi: 10.3390/md17030147
- Hughes K.A. and Stallwood B. 2005. Oil pollution in the Antarctic terrestrial environment. *Polarforschung* 75: 141–144. doi: 10013/epic.29948.d001
- Ibrahim S., Zulkharnain A., Zahri K.N.M., Lee G.L.Y., Convey P., Gomez-Fuentes C., Sabri S., Khalil K.A., Alias S.A., Gonzalez-Rocha G. and Ahmad S.A. 2020. Effect of heavy metals and other xenobiotics on biodegradation of waste canola oil by cold-adapted *Rhodococcus* sp. strain AQ5-07. *Revista Mexicana de Ingeniera Quimica* 19: 1041–1052. doi: 10.24275/rmiq/Bio917
- Jiang X., Hou S., Li Y., Pang H., Hua R., Paul M., Sharon S., An C., Michael H., Liu K. and Zhang W. 2018. Spatial variations of Pb, As, and Cu in surface snow along the transect from the Zhongshan Station to Dome A, East Antarctica. *Sciences in Cold and Arid Regions* 10: 0219–0231.
doi: 10.3724/SP.J.1226.2018.00219

- Kaamoush M., El-Agawany N., Salhin H.E. and El-Zeiny A. 2022. Monitoring effect of nickel, copper, and zinc on growth and photosynthetic pigments of *Spirulina platensis* with suitability investigation in Idku Lake. *Environmental Science and Pollution Research* 29: 78942–78959. doi: 10.1007/s11356-022-21328-1
- Kai E.X., Wan Johari W.L., Habib S., Yasid N.A., Ahmad S.A. and Shukor M.Y. 2020. The growth of the *Rhodococcus* sp. on diesel fuel under the effect of heavy metals and different concentrations of zinc. *Advances in Polar Science* 31: 132–136. doi: 10.13679/j.advps.2019.0043
- Kalhor A.X., Movafeghi A., Mohammadi-Nassab A.D., Abedi E. and Bahrami A. 2017. Potential of the green alga *Chlorella vulgaris* for biodegradation of crude oil hydrocarbons. *Marine Pollution Bulletin* 123: 286–290. doi: 10.1016/J.MARPOLBUL.2017.08.045
- Kan G., Miao J., Shi C. and Li G. 2006. Proteomic alterations of Antarctic ice microalga *Chlamydomonas* sp. under low temperature stress. *Journal of Integrative Plant Biology* 48: 965–970. doi: 10.1111/j.1744-7909.2006.00255.x
- Ke L., Luo L., Wang P., Luan T. and Tam N.F.Y. 2010. Effects of metals on biosorption and biodegradation of mixed polycyclic aromatic hydrocarbons by a freshwater green alga *Selenastrum capricornutum*. *Bioresource Technology* 101: 6950–6961. doi: 10.1016/J.BIORTECH.2010.04.011
- Kennicutt M.C. 1990. Oil spillage in Antarctica: Initial report of the National Science Foundation-sponsored Quick Response Team on the grounding of the *Bahia Paraiso*. *Environmental Science and Technology* 24: 620–624. doi: 10.1021/ES00075A601
- Kennicutt M.C., McDonald S.J., Sericano J.L., Boothe P., Oliver J., Safe S., Presley B.J., Liu H., Wolfe D., Wade T.L., Crockett A. and Bockus D. 1995. Human contamination of the marine environment-Arthur Harbor and McMurdo Sound, Antarctica. *Environmental Science and Technology* 29: 1279–1287. doi: 10.1021/es00005a600
- Kerry E. 1993. Bioremediation of experimental petroleum spills on mineral soils in the Vestfold Hills, Antarctica. *Polar Biology* 13: 163–170. doi: 10.1007/BF00238926
- Korashy H.M. and El-Kadi A.O.S. 2005. Regulatory mechanisms modulating the expression of cytochrome P450 1A1 gene by heavy metals. *Toxicological Sciences* 88: 39–51. doi: 10.1093/toxsci/kfi282
- Krzemińska I., Pawlik-Skowrońska B., Trzcńska M. and Tys J. 2014. Influence of photoperiods on the growth rate and biomass productivity of green microalgae. *Bioprocess and Biosystems Engineering* 37: 735–741. doi: 10.1007/s00449-013-1044-x
- Kumar A. and Bera S. 2020. Revisiting nitrogen utilization in algae: A review on the process of regulation and assimilation. *Bioresource Technology Reports* 12: 100584. doi: 10.1016/j.biteb.2020.100584
- Küpper H. 2017. Lead toxicity in plants. In: Sigel A., Sigel H. and Sigel R. (eds.) *Lead: Its Effects on Environment and Health*. De Gruyter: 491–500. doi: 10.1515/9783110434330-015

- Lichtenthaler H.K. and Buschmann C. 2001. Chlorophylls and carotenoids: Measurement and characterization by UV-VIS spectroscopy. *Current Protocols in Food Analytical Chemistry* F4.3.1-F4.3.8. doi: 10.1002/0471142913.faf0403s01
- Lim Z.S., Wong R.R., Wong C.Y., Zulkharnain A., Shaharuddin N.A. and Ahmad S.A. 2021. Bibliometric analysis of research on diesel pollution in Antarctica and a review on remediation techniques. *Applied Sciences* 11: 1123. doi: 10.3390/app11031123
- Lim Z.S., Wong C.Y., Ahmad S.A., Puasa N.A., Phang L.Y., Shaharuddin N.A., Merican F., Convey P., Zulkharnain A., Shaari H., Azmi A.A., Kok Y.Y. and Gomez-Fuentes C. 2023. Harnessing diesel-degrading potential of an Antarctic microalga from Greenwich Island and its physiological adaptation. *Biology* 12: 1142. doi: 10.3390/biology12081142
- Luo A., Wu Y.R., Xu Y., Kan J., Qiao J., Liang L., Huang T. and Hu Z. 2016. Characterization of a cytochrome P450 monooxygenase capable of high molecular weight PAHs oxidization from *Rhodococcus* sp. P14. *Process Biochemistry* 51: 2127–2133. doi: 10.1016/j.procbio.2016.07.024
- Łukowski A. and Dec D. 2021. Fractions of Ni, Pb, Cr, and their impact on enzyme activities of arable land cultivated by the simplified method. *Minerals* 11: 584. doi: 10.3390/min11060584
- Matcott J., Baylis S. and Clarke R.H. 2019. The influence of petroleum oil films on the feather structure of tropical and temperate seabird species. *Marine Pollution Bulletin* 138: 135–144. doi: 10.1016/j.marpolbul.2018.11.010
- McFarlin K.M., Perkins M.J., Field J.A. and Leigh M.B. 2018. Biodegradation of crude oil and Corexit 9500 in Arctic seawater. *Frontiers in Microbiology* 9: 1788. doi: 10.3389/FMICB.2018.01788/BIBTEX
- McGenity T.J., Folwell B.D., McKew B.A. and Sanni G.O. 2012. Marine crude-oil biodegradation: A central role for interspecies interactions. *Aquatic Biosystems* 8: 10. doi: 10.1186/2046-9063-8-10
- Monteiro C.M., Castro P.M.L. and Malcata F.X. 2012. Metal uptake by microalgae: Underlying mechanisms and practical applications. *Biotechnology Progress* 28: 299–311. doi: 10.1002/btpr.1504
- O'Brien P.P.Y. and Dixon P.S. 1976. The effects of oils and oil components on algae: A review. *British Phycological Journal* 11: 115–142. doi: 10.1080/00071617600650161
- Osipova S., Dudareva L., Bondarenko N., Nasarova A., Sokolova N., Obolkjna L., Glyzina O. and Timoshkin O. 2009. Temporal variation in fatty acid composition of *Ulothrix zonata* (Chlorophyta) from ice and benthic communities of Lake Baikal. *Phycologia* 48: 130–135. doi: 10.2216/08-49.1
- Pathak B., Gupta S. and Verma R. 2018. Biosorption and biodegradation of polycyclic aromatic hydrocarbons (PAHs) by microalgae. In: Crini G. and Lichtfouse E. (eds.) *Green Adsorbents for*

- Pollutant Removal*. Springer International Publishing: 215–247. doi: 10.1007/978-3-319-92111-2_7
- Pople A., Simpson R.D. and Cairns S.C. 1990. An incident of Southern Ocean oil pollution: Effects of a spillage of diesel fuel on the rocky shore of Macquarie Island (Sub-Antarctic). *Marine and Freshwater Research* 41: 603–620. doi: 10.1071/MF9900603
- Puasa N.A., Zulkharnain A., Verasoundarapandian G., Wong C.Y., Zahri K.N.M., Merican F., Shaharuddin N.A., Gomez-Fuentes C. and Ahmad S.A. 2021. Effects of diesel, heavy metals and plastics pollution on penguins in Antarctica: A review. *Animals* 11: 2505. doi: 10.3390/ani11092505
- Ramadass K., Megharaj M., Venkateswarlu K. and Naidu R. 2017. Toxicity of diesel water accommodated fraction toward microalgae, *Pseudokirchneriella subcapitata* and *Chlorella* sp. MM3. *Ecotoxicology and Environmental Safety* 142: 538–543. doi: 10.1016/j.ecoenv.2017.04.052
- Raven J.A., Evans M.C.W. and Korb R.E. 1999. The role of trace metals in photosynthetic electron transport in O₂-evolving organisms. *Photosynthesis Research* 60: 111–150. doi: 10.1023/A:1006282714942
- Raymond T., King C.K., Raymond B., Stark J.S. and Snape I. 2017. Oil pollution in Antarctica. In: Fingas M. (eds.) *Oil Spill Science and Technology*. Gulf Professional Publishing: 759–803. doi: 10.1016/B978-0-12-809413-6.00014-X
- Rezayian M., Niknam V. and Ebrahimzadeh H. 2019. Oxidative damage and antioxidative system in algae. *Toxicology Reports* 6: 1309–1313. doi: 10.1016/j.toxrep.2019.10.001
- Rocha G.S., Parrish C.C. and Espíndola E.L.G. 2021. Effects of copper on photosynthetic and physiological parameters of a freshwater microalga (Chlorophyceae). *Algal Research* 54: 102223. doi: 10.1016/j.algal.2021.102223
- Roslee A.F.A., Ahmad S.A., Gomez-Fuentes C., Shaharuddin N.A., Khalil K.A. and Zulkharnain A. 2021. Scientometric analysis of diesel pollutions in Antarctic territories: A review of causes and potential bioremediation approaches. *Sustainability* 13: 7064. doi: 10.3390/su13137064
- Sazhin S.S., Al Qubeissi M., Nasiri R., Gun'Ko V.M., Elwardany A.E., Lemoine F., Grisch F. and Heikal M.R. 2014. A multi-dimensional quasi-discrete model for the analysis of diesel fuel droplet heating and evaporation. *Fuel* 129: 238–266. doi: 10.1016/J.FUEL.2014.03.028
- Scarano G. and Morelli E. 2002. Characterization of cadmium- and lead- phytochelatin complexes formed in a marine microalga in response to metal exposure. *BioMetals* 15: 145–151. doi: 10.1023/A:1015288000218
- Sharma I. 2020. Bioremediation techniques for polluted environment: Concept, advantages, limitations, and prospects. In: Alfonso Murillo-Tovar M., Saldarriaga-Noreña H. and Saied A. (eds.) *Trace Metals in the Environment*. IntechOpen. doi: 10.5772/INTECHOPEN.90453

- Sieber M., Conway T.M., de Souza G.F., Hassler C.S., Ellwood M.J. and Vance D. 2019. High-resolution Cd isotope systematics in multiple zones of the Southern Ocean from the Antarctic Circumnavigation Expedition. *Earth and Planetary Science Letters* 527: 115799.
doi: 10.1016/j.epsl.2019.115799
- Sieber M., Conway T.M., de Souza G.F., Hassler C.S., Ellwood M.J. and Vance D. 2020. Cycling of zinc and its isotopes across multiple zones of the Southern Ocean: Insights from the Antarctic Circumnavigation Expedition. *Geochimica et Cosmochimica Acta* 268: 310–324.
doi: 10.1016/j.gca.2019.09.039
- Souza P.O., Ferreira L.R., Pires N.R.X., Filho P.J.S., Duarte F.A., Pereira C.M.P. and Mesko M.F. 2012. Algae of economic importance that accumulate cadmium and lead: A review. *Revista Brasileira de Farmacognosia Brazilian Journal of Pharmacognosy* 22: 825–837.
doi: 10.1590/S0102
- Souza-Kasprzyk J., de Castro Paiva T., Convey P., Schmauder Teixeira da Cunha L., Soares T.A., Zawierucha K., Costa E.S., Niedzielski P. and Machado Torres J.P. 2022. Influence of marine vertebrates on organic matter, phosphorus and other chemical element levels in Antarctic soils. *Polar Biology* 45: 1571–1580. doi: 10.1007/s00300-022-03091-8
- Sujetovienė G. 2014. Copper induced physiological changes and oxidative damage in lichen *Ramalina farinacea*. *Biologija* 60: 196–201. doi: 10.6001/BIOLOGIJA.V60I4.3039
- Sunda W.G. and Huntsman S.A. 1983. Effect of competitive interactions between manganese and copper on cellular manganese and growth in estuarine and oceanic species of the diatom *Thalassiosira*. *Limnology and Oceanography* 28: 924–934. doi: 10.4319/LO.1983.28.5.0924
- Sydney E.B., Novak A.C., de Carvalho J.C. and Soccol C.R. 2014. Respiriometric balance and carbon fixation of industrially important algae. In: Pandey A., Lee D.J., Chisti Y. and Soccol C.R. (eds.) *Biofuels from Algae*. Elsevier: 67–84. doi: 10.1016/B978-0-444-59558-4.00004-8
- Taniguchi S., Montone R.C., Bicego M.C., Colabuono F.I., Weber R.R. and Sericano J.L. 2009. Chlorinated pesticides, polychlorinated biphenyls and polycyclic aromatic hydrocarbons in the fat tissue of seabirds from King George Island, Antarctica. *Marine Pollution Bulletin* 58: 129–133.
doi: 10.1016/j.marpolbul.2008.09.026
- Tin T., Sovacool B.K., Blake D., Magill P., El Naggar S., Lidstrom S., Ishizawa K. and Berte J. 2010. Energy efficiency and renewable energy under extreme conditions: Case studies from Antarctica. *Renewable Energy* 35: 1715–1723. doi: 10.1016/j.renene.2009.10.020
- Tin T., Lamers M., Liggett D., Maher P.T. and Hughes K.A. 2014. Setting the scene: Human activities, environmental impacts and governance arrangements in Antarctica. In: Tin T., Liggett D., Maher P. and Lamers M. (eds.) *Antarctic Futures: Human Engagement with the Antarctic Environment*. Springer, Dordrecht: 1–24. doi: 10.1007/978-94-007-6582-5_1
- Touliabah H.E.S., El-Sheekh M.M., Ismail M.M. and El-Kassas H. 2022. A review of microalgae- and cyanobacteria-based biodegradation of organic pollutants. *Molecules* 27: 1141.

- doi: 10.3390/MOLECULES27031141
- Trevors J.T., Stratton G.W. and Gadd G.M. 1986. Cadmium transport, resistance, and toxicity in bacteria, algae, and fungi. *Canadian Journal of Microbiology* 32: 447–464.
doi: 10.1139/M86-085
- Tsuji N., Hirayanagi N., Okada M., Miyasaka H., Hirata K., Zenk M.H. and Miyamoto K. 2002. Enhancement of tolerance to heavy metals and oxidative stress in *Dunaliella tertiolecta* by Zn-induced phytochelatin synthesis. *Biochemical and Biophysical Research Communications* 293: 653–659. doi: 10.1016/S0006-291X(02)00265-6
- Tsuji N., Hirayanagi N., Iwabe O., Namba T., Tagawa M., Miyamoto S., Miyasaka H., Takagi M., Hirata K. and Miyamoto K. 2003. Regulation of phytochelatin synthesis by zinc and cadmium in marine green alga, *Dunaliella tertiolecta*. *Phytochemistry* 62: 453–459.
doi: 10.1016/S0031-9422(02)00559-9
- Tumeo M.A. and Larson M.K. 1994. Movement of fuel spills in the Ross Ice Shelf. *Antarctic Journal of United States*, 29.
- Vo P., Nhat H., Ngo H.H., Guo W.S., Chang S.W., Nguyen D.D., Nguyen P.D., Bui X.T., Zhang X.B. and Guo J.B. 2018. Can algae-based technologies be an affordable green process for biofuel production and wastewater remediation? *Bioresource Technology* 256: 491–501.
doi: 10.1016/j.biortech.2018.02.031
- Vymazal J. 1987. Toxicity and accumulation of cadmium with respect to algae and cyanobacteria: A review. *Toxicity Assessment* 2: 387–415. doi: 10.1002/tox.2540020403
- Wang Y., Ahmed Z., Feng W., Li C. and Song S. 2008. Physicochemical properties of exopolysaccharide produced by *Lactobacillus kefiranofaciens* ZW3 isolated from Tibet kefir. *International Journal of Biological Macromolecules* 43: 283–288.
doi: 10.1016/J.IJBIOMAC.2008.06.011
- Webb A.L., Hughes K.A., Grand M.M., Lohan M.C. and Peck L.S. 2020. Sources of elevated heavy metal concentrations in sediments and benthic marine invertebrates of the western Antarctic Peninsula. *Science of The Total Environment* 698: 134268.
doi: 10.1016/J.SCITOTENV.2019.134268
- Wong P.T.S. and Chau Y.K. 1990. Zinc toxicity to freshwater algae. *Toxicity Assessment* 5: 167–177.
doi: 10.1002/tox.2540050205
- Wong R.R., Lim Z.S., Shaharuddin N.A., Zulkharnain A., Gomez-Fuentes C. and Ahmad S.A. 2021. Diesel in Antarctica and a bibliometric study on its indigenous microorganisms as remediation agent. *International Journal of Environmental Research and Public Health* 18: 1512.
doi: 10.3390/IJERPH18041512
- Xu Q., Chu Z., Gao Y., Mei Y., Yang Z., Huang Y., Yang L., Xie Z. and Sun L. 2020. Levels, sources and influence mechanisms of heavy metal contamination in topsoils in Mirror Peninsula, East Antarctica. *Environmental Pollution* 257: 113552. doi: 10.1016/j.envpol.2019.113552

- Ye N., Han W., Toseland A., Wang Y. *et al.* 2022. The role of zinc in the adaptive evolution of polar phytoplankton. *Nature Ecology and Evolution* 6: 965–978. doi: 10.1038/s41559-022-01750-x
- Zahri K.N.M., Zulkharnain A., Gomez-Fuentes C., Sabri S. and Ahmad S.A. 2020. Effects of heavy metals on Antarctic bacterial cell growth kinetics and degradation of waste canola oil. *Journal of Environmental Biology* 41: 1433–1441. doi: 10.22438/JEB/41/6/MRN-1464
- Zahri K.N.M., Gomez-Fuentes C., Sabri S., Zulkharnain A., Khalil K.A., Lim S. and Ahmad S.A. 2021. Evaluation of heavy metal tolerance level of the Antarctic bacterial community in biodegradation of waste canola oil. *Sustainability* 13: 10749. doi: 10.3390/su131910749
- Zakaria N.N., Roslee A.F.A., Gomez-Fuentes C., Zulkharnain A., Abdulrasheed M., Sabri S., Ramirez-Moreno N., Calisto-Ulloa N. and Ahmad S.A. 2020. Kinetic studies of marine psychrotolerant microorganisms capable of degrading diesel in the presence of heavy metals. *Revista Mexicana de Ingeniera Quimica* 19: 1375–1388. doi: 10.24275/rmiq/Bio1072
- Zakaria N.N., Convey P., Gomez-Fuentes C., Zulkharnain A., Sabri S., Shaharuddin N.A. and Ahmad S.A. 2021. Oil bioremediation in the marine environment of Antarctica: A review and bibliometric keyword cluster analysis. *Microorganisms* 9: 419. doi: 10.3390/MICROORGANISMS9020419
- Zhao Y., Song X., Yu L., Han B., Li T. and Yu X. 2019. Influence of cadmium stress on the lipid production and cadmium bioresorption by *Monoraphidium* sp. QLY-1. *Energy Conversion and Management* 188: 76–85. doi: 10.1016/J.ENCONMAN.2019.03.041
- Zheng S., Guo J., Cheng F., Gao Z., Du L., Meng C., Li S. and Zhang X. 2022. Cytochrome P450s in algae: Bioactive natural product biosynthesis and light-driven bioproduction. *Acta Pharmaceutica Sinica B* 12: 2832–2844. doi: 10.1016/J.APSB.2022.01.013

Received 4 September 2024

Accepted 7 November 2024

Table 1.

Specific growth rate (μ) of *Tritostichococcus* sp. WCY_AQ5_1 under exposure to different concentrations of one of four heavy metals over a 7 d culture period.

Treatment (ppm)	Specific growth rate (μ), d ⁻¹			
	Zn	Pb	Cu	Cd
0	0.135 ± 0.022	0.189 ± 0.013	0.115 ± 0.003	0.115 ± 0.003
1	0.144 ± 0.017	0.221 ± 0.044	0.116 ± 0.006	0.106 ± 0.009
2	0.151 ± 0.011	0.194 ± 0.003	0.116 ± 0.007	0.092 ± 0.011
4	0.141 ± 0.010	0.209 ± 0.012	0.121 ± 0.006	0.107 ± 0.011
8	0.149 ± 0.003	0.202 ± 0.025	0.108 ± 0.001	0.117 ± 0.021
16	0.033 ± 0.026*	0.128 ± 0.018*	0.018 ± 0.01*	0.073 ± 0.011

Data shown are mean values ± standard deviation of three replicates. Asterisks (*) indicate significant differences at $p < 0.05$.




Fig. 1. Growth of *Tritostichococcus* sp. WCY_AQ5_1 over 7 d exposure to (A) Zn, (B) Pb, (C) Cu, and (D) Cd at various concentrations. Data shown are mean values of three replicates with error bars indicating standard deviation.




Fig. 2. Photosynthetic pigment content of *Tritostichoecoccus* sp. WCY_AQ5_1 on day 7 of exposure to different concentrations of (A) Zn, (B) Pb, (C) Cu and (D) Cd. Data shown are mean values of three replicates with error bars indicating standard deviation. Asterisks in blue (*), brown (*), orange (*) and hashtag (#) denote significant differences at $p < 0.05$ for chl-*a*, chl-*b*, car and car/chl-*a* ratio, respectively.




Fig. 3. Growth of *Tritostichococcus* sp. WCY_AQ5_1 on day 7 under (A) Zn, (B) Pb, (C) Cu, and (D) Cd treatment at various concentrations with and without diesel co-treatment. Data shown are mean values of three replicates with error bars indicating standard deviation. Asterisks denote significant differences at $p < 0.05$.




Fig. 4. Biodegradation efficiency of diesel and growth of *Tritostichococcus* sp. WCY_AQ_1 in BBM that contained diesel and (A) Zn, (B) Pb, (C) Cu, or (D) Cd, measured on day 7. Data shown are mean values of three replicates with error bars indicating standard deviation. Uppercase letters (A, B, C) and lowercase letters (a, b, c) denote significant differences at $p < 0.05$ for chl-*a* and BE, respectively, between treatment groups for each heavy metal.




Fig. 5. Photosynthetic pigment content of *Tritostichococcus* sp. WCY_AQ5_1 on day 7 in BBM that contained 1% v/v diesel and different concentrations of (A) Zn, (B) Pb, (C) Cu, and (D) Cd. Data shown are mean values of three replicates with error bars indicating standard deviation. Uppercase letters (A, B, C), lowercase letters (a, b, c), lowercase letters with asterisk (a*, b*, c*), lowercase letters in green (a, b, c) and pink (a, b, c) denote significant differences at $p < 0.05$ in chl-*a*, chl-*b* and car contents, and chl-*b*/chl-*a* and car/chl-*a* ratios, respectively, between treatment groups for each heavy metal.




Fig. 6. Reactive oxygen species (ROS) levels (A–D) and lipid peroxidation (LP) levels (E–H) in *Tritostichococcus* sp. WCY_AQ5_1 on day 7 in culture in BBM containing 1% v/v initial diesel concentration in combination with different concentrations of Zn, Pb, Cu, or Cd. Data shown are mean values of three replicates with error bars indicating standard deviation. Lowercase letters (a, b, c, d, e) denote significant difference at $p < 0.05$ in ROS and LP between treatment groups for each heavy metal.