

RAPID COMMUNICATION

Monitoring the occurrence of bacteria in stored cabbage heads

Aleš Eichmeier¹, Eliška Peňázová², Jakub Pečenka¹, Jana Čechová¹, Robert Pokluda², Dorota Tekielska^{3*}, Miroslav Baránek¹

¹ Mendeleum – Institute of Genetics and Plant Breeding, Mendel University in Brno, Faculty of Horticulture, Valtická 337, 691 44 Lednice, Czech Republic

² Department of Vegetable Growing and Floriculture, Faculty of Horticulture, Mendel University in Brno, Valtická 337, 691 44 Lednice, Czech Republic

³ Department of Plant Protection, Faculty of Biotechnology and Horticulture, University of Agriculture in Krakow, 29 Listopada 54, 31-425 Krakow, Poland

Vol. 57, No. 1: 56–61, 2017

DOI: 10.1515/jppr-2017-0008

Received: December 5, 2016

Accepted: January 23, 2017

*Corresponding address:
d.tekielska@gmail.com

Abstract

Twenty-six cabbage heads stored under typical conditions in a storage hall in Moravia, Czech Republic, were tested for the presence of bacteria by the method of isolation from three different parts of the cabbage heads. Isolations were carried out from stalks, inner and superficial leaves. Two samplings were done; in November 2015 and February 2016. Bacterial cultures were sequenced in the part of 16S rRNA region; bacteria were identified according to the sequences obtained. The most prevalent bacteria were of the genus *Pseudomonas*. Genera: *Klebsiella*, *Erwinia*, *Pantoea*, *Bacillus* were also identified. The results provided an interesting insight into the bacterial spectrum in stored cabbage heads and their dynamics during storage. The nucleotide sequences which were found were saved in GenBank/NCBI under accession numbers KX160104-KX160145.

Key words: *Bacillus*, bacteria, cabbage, identification, isolation, MPA medium, *Pseudomonas*, sequencing, 16S rRNA

Introduction

Bacterial soft rot pathogens may cause serious losses in stored cabbage heads (*Brassica oleracea* var. *capitata* L.). Vegetables may already be infected at harvest even though they do not show visible symptoms (Bhat *et al.* 2012). Due to the ability to cause latent infection, the appearance of symptoms is strongly dependent on environmental conditions (Bretschneider *et al.* 1989; Czajkowski *et al.* 2011). However, postharvest bacterial soft rot losses caused by various species of *Bacillus*, *Pseudomonas* and *Erwinia* have been estimated to vary between 15–30% of the harvested crop (Agrios 2006). Bacteria which infect cruciferous plants are primarily gram-negative. The most frequently identified genera are *Pseudomonas* (Mauzey *et al.* 2015), *Erwinia* (Togashi *et al.* 2001), *Pectobacterium* (Nazerian *et al.* 2011), *Xanthomonas* (Roberts *et al.* 1999; Eichmeier

et al. 2015) and *Bacillus* (Agrios 2006). The bacteria mainly attack the fleshy storage organs of their hosts (Bhat *et al.* 2010). Soft rot pathogens are transmitted by contact with infected plants or by their persistence on postharvest residues (Davidsson *et al.* 2013). Infection can also be spread by insects (Nadarasah *et al.* 2011).

Control of the disease is not always very effective, however, sanitary practices in production, storage and processing can slow the spread of the disease and protect yields (Czajkowski *et al.* 2011). Furthermore, dispersal of bacteria can also happen via usage of surface water for irrigation, aerosols generated by rain, movement of bacteria in soil water or mechanically via contaminated agricultural equipment (Perombelon and Kelman 1980; Czajkowski *et al.* 2011).

This work is focused on the determination of bacterial communities present in different parts of cabbage heads and their dynamics during cold storage.

Materials and Methods

Sources of materials and samplings

The bacterial isolates originated from naturally infected plants of cabbage *B. oleracea* var. *capitata* cv. Fundaxy, the cultivar of white cabbage suitable for storage. Cabbage heads were harvested in October 2015 in Otice (GPS coordinates: 49.91683940000001, 17.86982660000001) and they were stored at 1–2°C and high humidity (95%) in a storage hall located in Otice. Two samplings were done: in November 2015 and February 2016. At both samplings, 13 cabbage heads were analysed.

Isolation of bacteria from cabbage heads

From each head, three samples from different locations within the cabbage head were taken as follows: A – cabbage stalk, B – inner leaves and C – superficial leaves. The heads were cut with a knife and scalpel. Approximately 4 × 4 mm pieces were disinfected in 2% sodium hypochlorite solution and washed 2× in sterile distilled water. The pieces from parts A, B and C were placed on Petri dishes. Samples were cultivated on MPA (meat-peptone agar, Sigma-Aldrich, St. Louis, USA) on Petri dishes at 25°C in the dark. Most of the cabbage heads were without symptoms but some of them showed symptoms e.g. black spots on the superficial leaves – C. Parts A and B were entirely without symptoms (Tables 1 and 2).

PCR reaction and sequencing

The DNA from cultures colonising each Petri dish was isolated from approximately 5 mg of bacterial culture according to Roohie and Umesha (2012). Target amplicon, the part of the 16S rRNA, V3 and V4 region (Klindworth *et al.* 2013) was amplified with GoTaq® G2 Flexi kit (Promega, Madison, USA). The PCR products corresponding to the expected size were gel-purified using NucleoSpin Gel and PCR Clean-up (Macherey-Nagel, Düren, Germany) and subjected to nucleotide sequencing as described by Eichmeier *et al.* (2010). The obtained nucleotide sequences were analysed using CLC Genomics Workbench 6.0 (CLC Bio, Aarhus, Denmark). The genus of bacteria was determined with similarities greater than 90% in GenBank/NCBI as recommended by Klindworth *et al.* (2013). The species level was based on 100% nucleotide identity.

Results

In the first sampling, bacterial colonies grew on 56% of the Petri dishes (Table 1). Two bacterial colonies (12C and 13A) were not identified because of the very high similarity with undetermined bacteria in GenBank/NCBI. Most of the identified colonies belonged to the genus *Pseudomonas*. Bacterial spectrum isolated from the first sampling is shown in Fig. 1 (Bacterial spectrum from the 1st sampling).

Bacterial cultures in the second sampling were obtained on 51% of Petri dishes (Table 2). The rates of bacteria isolated from the second sampling are shown in Fig. 2 (Bacterial spectrum from the second sampling).

Six genera of bacteria in 26 stored cabbage heads were detected based on DNA sequencing of cultivated colonies. In the first sampling five genera were present: *Erwinia*, unclassified *Gammaproteobacterium*, *Klebsiella*, *Pantoea* and *Pseudomonas*. In the second sampling only *Pseudomonas* and *Bacillus* colonies occurred.

Discussion

Plants and their products gradually lose their defence mechanisms against invading microorganisms after harvest. Therefore they are generally more susceptible to infection by natural pathogens or even to attack by bacteria that normally do not cause diseases of plants growing in the field (Liao and Wells 1987). King and Bolin (1989) indicated the increase of microbial populations of lettuce under storage conditions which makes our results surprising since we found that the spectrum of identified bacterial pathogens was lower in the second sampling than in the first one. This is one of the first studies which focused on identification of bacteria in different parts (A, B, C) of cabbage heads. A comparison of symptomatic and asymptomatic heads was carried out. We detected only *Pseudomonas* spp. in the first sampling in the asymptomatic heads. In the second sampling we also detected *Bacillus* spp. (5A, 7B) (Tables 1 and 2). In the symptomatic heads, we detected *Klebsiella*, *Erwinia*, *Pantoea* and *Pseudomonas* species in the first sampling and in the second sampling only *Bacillus* and *Pseudomonas* species. The occurrence of *Pseudomonas* spp. in both sampling periods and in both symptomatic and asymptomatic heads was expected because bacteria belonging to this genus are able to grow at low temperatures. Brocklehurst and Lund (1981) isolated five strains of *Pseudomonas* sp. from cabbage stored at 1°C.

No *Bacillus* spp. were observed in the first sampling. The presence of *Bacillus* spp. in the second sampling

Table 1. List of detected bacteria in cabbage heads in storage hall: A – stalk, B – inner leaves, C – superficial leaves, 1st sampling

Isolate	S/A*	Detected bacteria	blastN [%]	Acces. Number [GenBank]
1A	A	<i>Pseudomonas</i> sp.	95	KX160117
4A	A	<i>Pseudomonas</i> sp.	98	KX160118
4C	A	<i>Pseudomonas</i> sp.	98	KX160119
5C	A	<i>Pseudomonas</i> sp.	97	KX160120
6A	A	<i>Pseudomonas fluorescens</i>	100	KX160121
6C	A	<i>Pseudomonas</i> sp.	99	KX160122
8A	S	<i>Pseudomonas</i> sp.	96	KX160123
8B	S	<i>Pseudomonas</i> sp.	97	KX160124
8C	S	<i>Klebsiella</i> sp.	92	KX160125
9A	S	<i>Gammaproteobacterium</i> sp.	92	KX160126
9B	S	<i>Pseudomonas</i> sp.	99	KX160127
9C	S	<i>Pseudomonas</i> sp.	99	KX160128
10A	S	<i>Pseudomonas</i> sp.	93	KX160129
10B	S	<i>Pseudomonas</i> sp.	96	KX160130
10C	S	<i>Pseudomonas</i> sp.	98	KX160131
11A	S	<i>Erwinia</i> sp.	99	KX160104
11C	S	<i>Pantoea</i> sp.	99	KX160105
12A	S	<i>Pseudomonas</i> sp.	99	KX160106
12C	S	not identified	93	KX160107
13A	S	not identified	94	KX160108
13B	S	<i>Gammaproteobacterium</i> sp.	91	KX160109
13C	S	<i>Pseudomonas</i> sp.	92	KX160110

*S/A – symptomatic/asymptomatic heads




Fig. 1. Bacterial spectrum from the 1st sampling. The part of the circle which emerges from the unit belongs to asymptomatic cabbage heads (*Pseudomonas* spp.). The rest of the circle belongs to symptomatic heads.

was not surprising because spores of genus *Bacillus* have the ability to germinate and grow at refrigeration temperatures in different vegetable substrates (Nicholson *et al.* 2000; Valero *et al.* 2003).

Warth (1978) reported that the *Bacillus* spp. grow at 5°C as a minimal temperature. Valero *et al.* (2003) published results about *Bacillus* spp. growth at low temperatures and indicated that the pH strongly affects the

Table 2. List of detected bacteria in cabbage heads in storage hall: A – stalk, B – inner leaves, C – superficial leaves, 2nd sampling

Isolate	S/A*	Detected bacteria	blastN [%]	Acces. Number [GenBank]
2C	A	<i>Pseudomonas</i> sp.	98	KX160132
3C	A	<i>Pseudomonas</i> sp.	99	KX160133
4B	A	<i>Pseudomonas</i> sp.	100	KX160134
4C	A	<i>Pseudomonas</i> sp.	99	KX160135
5A	A	<i>Bacillus</i> sp.	87	KX160136
6A	A	<i>Pseudomonas</i> sp.	98	KX160137
6B	A	<i>Pseudomonas</i> sp.	94	KX160138
6C	A	<i>Pseudomonas</i> sp.	99	KX160139
7B	A	<i>Bacillus</i> sp.	96	KX160140
7C	A	<i>Pseudomonas</i> sp.	99	KX160141
8B	S	not identified	99	KX160142
8C	S	<i>Pseudomonas fluorescens</i>	100	KX160143
9C	S	not identified	99	KX160144
10C	S	<i>Pseudomonas</i> sp.	99	KX160145
11B	S	<i>Pseudomonas</i> sp.	99	KX160111
11C	S	<i>Pseudomonas</i> sp.	100	KX160112
12A	S	<i>Bacillus</i> sp.	99	KX160113
12B	S	<i>Bacillus</i> sp.	99	KX160114
12C	S	<i>Bacillus</i> sp.	99	KX160115
13A	S	<i>Pseudomonas</i> sp.	100	KX160116

*S/A – symptomatic/asymptomatic heads




Fig. 2. Bacterial spectrum from the 2nd sampling. The part of the circle which emerges from the unit belongs to asymptomatic cabbage heads (*Pseudomonas* spp. and *Bacillus* spp.). The rest of the circle belongs to symptomatic heads.

growth of colonies. The vegetable substrates provide suitable pH conditions. Our study confirms the presence of *Bacillus* spp. in cabbage stalks (Table 2) also at low temperatures. *Pseudomonas* spp. were present in the first sampling in 36% and in the second one in 33% of isolations, throughout the whole cabbage head. *Pseudomonas* spp. is able to grow at 4°C as the lowest temperature (De Jonghe *et al.* 2011). Strong antagonistic

behaviour of *Pseudomonas* spp. (Afsharmanesh *et al.* 2010) and *Bacillus* spp. against other bacteria (Zhao *et al.* 2013, 2014) could explain why the colonies of other genera declined with time inside the cabbage heads. *Bacillus* species are considered to be a potential biological control against postharvest diseases of vegetables and fruits, due to their antagonistic behaviour (Zhao *et al.* 2013; Arzanlou *et al.* 2016).

Conclusions

In the current study, the bacterial spectrum of genera *Pseudomonas*, *Klebsiella*, *Erwinia*, *Pantoea* and *Bacillus* was detected. Bacteria were detected in all of the examined parts of cabbage heads: stalk, inner leaves and superficial leaves.

Differences in bacterial spectrum present in the stored cabbage heads, depending on the time of sampling term were observed.

Development of *Pseudomonas* and *Bacillus* bacterial species at the later period of storage and dieback of other genera colonies may be due to the antagonistic behaviour of *Pseudomonas* and *Bacillus* spp.

The study proves the change of the bacterial spectrum through the whole cabbage heads under storage conditions even when the heads did not show significant and visible symptoms. Our results also show that the diversity of bacterial communities decreased during cold storage. Based on the obtained results future studies will focus on confirming the antagonistic effects of *Pseudomonas* and *Bacillus* spp. against the other genera.

Acknowledgements

This research was supported by The Ministry of Agriculture of the Czech Republic, Project No. QJ1510088. Access to computing and storage facilities owned by involved parties and projects contributing to the National Grid Infrastructure MetaCentrum, provided under the programme “Projects of Large Infrastructure for Research, Development and Innovations” (LM2010005), is greatly appreciated.

References

- Afsharmanesh H., Ahmadzadeh M., Javan-Nikkhah M., Behboudi K. 2010. Characterization of the antagonistic activity of a new indigenous strain of *Pseudomonas fluorescens* isolated from onion rhizosphere. *Journal of Plant Pathology* 92 (1):187–194.
- Agrios G.N. 2006. *Bacterial Soft Rots*. 5th Edn. Academic Press, San Diego, USA.
- Arzanlou M., Mousavi S., Bakhshi M., Khakvar R., Bandehagh A. 2016. Inhibitory effects of antagonistic bacteria inhabiting the rhizosphere of the sugarbeet plants, on *Cercospora beticola* Sacc., the causal agent of Cercospora leaf spot disease on sugarbeet. *Journal of Plant Protection Research* 56 (1): 6–14.
- Bhat K.A., Bhat N.A., Mohiddin F.A., Mir S.A., Mir M.R. 2012. Management of post-harvest *Pectobacterium* soft rot of cabbage (*Brassica oleracea* var *capitata* L.) by biocides and packing material. *African Journal of Agricultural Research* 7 (28): 4066–4074.
- Bhat K.A., Masood S.D., Bhat N.A., Bhat M.A., Razvi S.M., Mir M.R., Habib M. 2010. Current status of postharvest soft rot in vegetables. *Asian Journal of Plant Sciences* 9 (4): 200–208.
- Bretschneider K.E., Gonella M.P., Robeson D.J. 1989. A comparative light and electron microscopical study of compatible and incompatible interactions between *Xanthomonas campestris* pv. *campestris* and cabbage (*Brassica oleracea*). *Physiological and Molecular Plant Pathology* 34 (4): 285–297.
- Brocklehurst T., Lund B.M. 1981. Properties of pseudomonads causing spoilage of vegetables stored at low temperature. *Journal of Applied Bacteriology* 50 (2): 259–266.
- Czajkowski R., Pérombelon M.C.M., Van Veen J.A., Van der Wolf J.M. 2011. Control of blackleg and tuber soft rot of potato caused by *Pectobacterium* and *Dickeya* species: a review. *Plant Pathology* 60 (6): 999–1013.
- Davidsson P.R., Kariola T., Niemi O., Palva E.T. 2013. Pathogenicity of and plant immunity to soft rot pectobacteria. *Frontiers in Plant Science* 4: 191.
- De Jonghe V., Coorevits A., Van Hoorde K., Messens W., Van Landschoot A., De Vos P., Heyndrickx M. 2011. Influence of storage conditions on the growth of *Pseudomonas* species in refrigerated raw milk. *Applied and Environmental Microbiology* 77 (2): 460–470.
- Eichmeier A., Baránek M., Pidra M. 2010. Analysis of genetic diversity and phylogeny of partial coat protein domain in Czech and Italian GFLV isolates. *Plant Protection Science* 46 (8): 145–148.
- Eichmeier A., Čechová J., Peňázová E. 2015. Genetic diversity of partial hrpF and Zur genes in populations of *Xanthomonas campestris* pv. *campestris* in *Brassica oleracea* convar. *capitata* in the Czech Republic. *Acta Horticulturae* 1105: 180–188.
- King A.D., Bolin H.R. 1989. Physiological and microbiological storage stability of minimally processed fruits and vegetables. *Food Technology* 43 (2): 132–139.
- Klindworth A., Pruesse E., Schweer T., Peplies J., Quast C., Horn M., Glöckner F.O. 2013. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Research*. Available on: <https://doi.org/10.1093/nar/gks808> [Accessed: September 4, 2016]
- Liao C.H., Wells J.M. 1987. Association of pectolytic strains of *Xanthomonas campestris* with soft rots of fruits and vegetables at retail markets. *Phytopathology* 77: 418–422.
- Mauzey S.J., Koike S.T., Bull C.T. 2015. First report of bacterial blight of cabbage (*Brassica oleracea* var. *capitata*) caused by *Pseudomonas cannabina* pv. *alisalensis* in California. *Journal of General Plant Pathology* 81: 331–350.
- Nadarasah G., Stavrinides J. 2011. Insects as alternative hosts for phytopathogenic bacteria. *FEMS Microbiology Reviews* 35: 555–575.
- Nazerian E., Sijam K., Mior Ahmad Z.A., Vadamalai G. 2011. First report of cabbage soft rot caused by *Pectobacterium carotovorum* subsp. *carotovorum* in Malaysia. *Plant Disease* 95 (4): 491–491.
- Nicholson W.L., Munakata N., Horneck G., Melosh H.J., Setlow P. 2000. Resistance of *Bacillus* endospores to extreme terrestrial and extraterrestrial environments. *Microbiology and Molecular Biology Reviews* 64 (3): 548–572.
- Perombelon M.C.M., Kelman A. 1980. Ecology of the soft rot erwinias. *Annual Review of Phytopathology* 18: 361–387.
- Roberts S.J., Hiltunen L.H., Hunter P.J., Brough J. 1999. Transmission from seed to seedling and secondary spread of *Xanthomonas campestris* pv. *campestris* in *Brassica* transplants: effects of dose and watering regime. *European Journal of Plant Pathology* 105: 879–889.
- Roohie R.K., Umesha S. 2012. Development of multiplex PCR for the specific detection of *Xanthomonas campestris* pv. *campestris* in cabbage and correlation with disease incidence. *Journal of Plant Pathology and Microbiology* 3: 1–9.
- Togashi J., Koushi U.E.D.A., Namai T. 2001. Overwintering of *Erwinia carotovora* subsp. *carotovora* in diseased tissues in soil and its role as inoculum for soft rot of Chinese cabbage (*Brassica campestris*, *Pekinensis* group). *Journal of General Plant Pathology* 67 (1): 45–50.

- Valero M., Fernandez P.S., Salmeron M.C. 2003. Influence of pH and temperature on growth of *Bacillus cereus* in vegetable substrates. *International Journal of Food Microbiology* 82 (1): 71–79.
- Warth A.D. 1978. Relationship between the heat resistance of spores and the optimum and maximum growth temperatures of *Bacillus* species. *Journal of Bacteriology* 134 (3): 699–705.
- Zhao Y., Li P., Huang K., Wang Y., Hu H., Sun Y. 2013. Control of postharvest soft rot caused by *Erwinia carotovora* of vegetables by a strain of *Bacillus amyloliquefaciens* and its potential modes of action. *World Journal of Microbiology and Biotechnology* 29 (3): 411–420.
- Zhao Y., Selvaraj J.N., Xing F., Zhou L., Wang Y., Song H., Liu Y. 2014. Antagonistic action of *Bacillus subtilis* strain SG6 on *Fusarium graminearum*. *PLoS One* 9 (3): e92486.