

RAPID COMMUNICATION

Effects of bacterial populations, temperature and exogenous hydrogen peroxide on the induction of the hypersensitive response in *Nicotiana tabacum* against *Xanthomonas perforans*

Ali Safaie Farahani, Seyyed Mohsen Taghavi*

Department of Plant Protection, College of Agriculture, Shiraz University, Shiraz, Iran

Vol. 57, No. 2: 201–204, 2017

DOI: 10.1515/jppr-2017-0019

Received: March 12, 2017

Accepted: May 24, 2017

*Corresponding address:
mtaghavi@shirazu.ac.ir

Abstract

The objective of this study was to investigate the effects of inoculum concentration, plant post-inoculation incubation temperature and exogenous hydrogen peroxide (H_2O_2) on the induction of the hypersensitive response (HR) in *Nicotiana tabacum* against *Xanthomonas perforans*. Inoculation of leaves with *X. perforans* at a concentration of 10^8 CFU · ml⁻¹ and incubation of plants at 30°C resulted in the strongest HR elicitation. Furthermore, an exogenous supply of H_2O_2 accelerated *X. perforans*-induced HR, whereas *in planta* H_2O_2 removal by application of catalase led to a delay in HR development. Our data suggest that H_2O_2 has an important role in HR of *N. tabacum* against *X. perforans*.

Key words: catalase, hydrogen peroxide, hypersensitive response, *Nicotiana tabacum*, *Xanthomonas perforans*

Plants are constantly exposed to various biotic and abiotic stresses during their life cycle. Among biotic challenges, diseases caused by fungi, bacteria, viruses and nematodes result in serious yield losses worldwide. In general, only a few pathogens have the ability to cause disease in a certain plant species. The resistance exhibited by an entire plant species to all genetic variants of a non-adapted pathogen species is referred to as non-host resistance. It is a durable and broad-spectrum resistance against numerous pathogens (Fan and Doerner 2012; Senthil-Kumar and Mysore 2013). Non-host resistance usually leads to the induction of the hypersensitive response (HR) at the infection site (Senthil-Kumar and Mysore 2013). The HR as the induction of quick necrotic lesions within 24–48 h after inoculation of tobacco leaves by incompatible plant pathogenic bacteria was first reported almost 50 years ago (Klement *et al.* 1964). Cell death rapidly encloses the site of infection and makes it almost impossible to develop biotrophic pathogens. One of the earliest responses of plants to pathogen invasion is the production of reactive oxygen species (ROS) including singlet oxygen (O_2), superoxide (O_2^-), hydrogen peroxide (H_2O_2) and

hydroxyl radical ($HO\cdot$) (Bolwell and Daudi 2009). Reactive oxygen species can restrict the pathogen through various mechanisms such as a direct effect on the invading pathogen (Levine *et al.* 1994), strengthening cell walls via oxidative cross-linking (Bradley *et al.* 1992), induction of systemic acquired resistance (Alvarez *et al.* 1998) and HR (Lamb and Dixon 1997). Accumulation of ROS in several non-host pathosystems such as barley/*Blumeria graminis* f. sp. *tritici* (Hückelhoven *et al.* 2001), cowpea/*Erysiphe cichoracearum* (Mellersh *et al.* 2002) and pepper/*Blumeria graminis* f. sp. *tritici* (Hao *et al.* 2011) has been reported. On the other hand, plants employ scavenging enzymes such as superoxide dismutase, ascorbate peroxidase and catalase for ROS detoxification (Mittler *et al.* 2004). The induction of scavenging enzymes in a number of non-host interactions such as pepper/*Xanthomonas campestris* pv. *campestris* (Kwak *et al.* 2009), broad bean/*Puccinia striiformis* f. sp. *tritici* (Cheng *et al.* 2012), bean/*X. hortorum* pv. *pelargonii* (Safaie Farahani and Taghavi 2015) and mung bean/*X. hortorum* pv. *pelargonii* (Safaie Farahani and Taghavi 2016) has been demonstrated. *Xanthomonas perforans* is the causal agent of bacterial

Table 1. The percentage of tobacco leaves showing hypersensitive response (HR) symptoms in different concentrations of bacterial inoculum and temperatures at 48 hpi (hours post-inoculation)

Temperature [°C]	Bacterial population		
	10 ⁶ CFU · ml ⁻¹	10 ⁷ CFU · ml ⁻¹	10 ⁸ CFU · ml ⁻¹
20	0±0	17±2	50±4
25	25±1	50±5	83±4
30	42±2	67±5	100±0

Table 2. The percentage of tobacco leaves showing hypersensitive response (HR) symptoms after exogenous application of H₂O₂ at different time intervals

Time intervals [hpi]	Treatment			
	control	H ₂ O ₂ at 4/6 hpi	H ₂ O ₂ at 6/8 hpi	H ₂ O ₂ at 8/10 hpi
12	33±2	73±3	60±6	40±2
24	80±4	100±0	100±0	100±0

hpi – hours post-inoculation

spot of tomato, an important disease worldwide (Jones *et al.* 2004). Optimum experimental situations and the role of ROS in non-host resistance of *Nicotiana tabacum* against *X. perforans* remain unclear. The goal of this study was to examine the role of temperature, bacterial populations and exogenous hydrogen peroxide in *X. perforans*-induced HR on tobacco leaves.

Xanthomonas perforans Tom801 (Osdaghi *et al.* 2016) was used in this study. To determine the role of bacterial populations and plant post-inoculation incubation temperature during *X. perforans*-induced HR, the inoculum at three concentrations (10⁶ CFU · ml⁻¹; 10⁷ CFU · ml⁻¹ or 10⁸ CFU · ml⁻¹) of bacterial suspension was separately infiltrated in fully expanded leaves of eight-week-old *N. tabacum* plants by needleless syringes. Following infiltration, the plants were incubated at 20°C, 25°C or 30°C. The percentage of leaves showing HR symptoms were recorded at 48 hours post inoculation (hpi). Twelve leaves from three plants were used for each treatment. To investigate the effects of H₂O₂ exogenous application on *X. perforans*-induced HR, the pathogen inoculum at a concentration of 10⁸ CFU · ml⁻¹ was infiltrated in fully expanded leaves. Afterwards, H₂O₂ at a concentration of 0.1 mM was sprayed twice (to maintain the high concentration of H₂O₂) to the inoculated leaf area at 4/6, 6/8 and 8/10 hpi. Control plants were treated with water instead of H₂O₂. The plants were incubated at 30°C. Necrosis symptoms were checked at 12 and 24 hpi. Fifteen leaves from four plants were used for each treatment. To remove *in planta* H₂O₂, an exogenous catalase was used. First, a bacterial suspension at a concentration of 10⁸ CFU · ml⁻¹ was infiltrated in fully expanded

leaves. In the next step, the catalase at a concentration of 2,000 U · ml⁻¹ was infiltrated twice to the inoculated leaf area at 4/6, 6/8 and 8/10 hpi. The plants were incubated at 30°C. Hypersensitive response symptoms were recorded at 24 and 48 hpi. Fifteen leaves from four plants were used for each treatment.

The optimal inoculum concentration and plant post-inoculation incubation temperature for *X. perforans*-induced HR were determined in this study. A high concentration of bacterial inoculum (10⁸ CFU · ml⁻¹) led to more induction of the HR than lower concentrations (10⁶ CFU · ml⁻¹ and 10⁷ CFU · ml⁻¹). Additionally, the plants that were incubated at 30°C showed more induction of the HR than those which were kept at 20°C and 25°C. Interaction of bacterial populations and temperature was also effective in HR induction. The greatest induction of the HR was found at 10⁸ CFU · ml⁻¹/30°C. On the other hand, no HR symptoms were observed at 10⁶ CFU · ml⁻¹/20°C (Table 1). There are several reports indicating different effects of inoculum concentration and plant post-inoculation incubation temperature on HR induction and non-host resistance (Jahnen and Hahlbrock 1988; Budde and Ullrich 2000; Li *et al.* 2015). This suggests that the role of these factors might depend on plant species, pathogen and their interaction. However, how the bacterial populations and temperature affect *X. perforans*-induced HR is still worth studying. Exogenous application of H₂O₂ led to faster and greater induction of the HR than the control. Furthermore, earlier exogenous H₂O₂ application (4/6 hpi) resulted in more HR elicitation than other H₂O₂ treatments (6/8 hpi and 8/10 hpi) at 12 hpi. There was no difference in terms of HR symptoms between the

Table 3. The percentage of tobacco leaves showing hypersensitive response (HR) symptoms after exogenous application of catalase at different time intervals

Time intervals [hpi]	Treatment			
	control	catalase at 4/6 hpi	catalase at 6/8 hpi	catalase at 8/10 hpi
24	80±7	40±4	53±3	67±6
48	100±0	53±1	73±6	93±4

hpi – hours post-inoculation

three H₂O₂ treatments at 24 hpi (Table 2). The leaves treated with catalase showed fewer HR symptoms than the control at 24 and 48 hpi. In addition, earlier catalase application (4/6 hpi) led to a greater reduction in HR induction than other catalase treatments (6/8 hpi and 8/10 hpi) at 24 and 48 hpi (Table 3). Our results are in agreement with a previous study by Li *et al.* (2015) that showed that an exogenous supply of H₂O₂ accelerates *Xanthomonas oryzae* pv. *oryzae*-induced HR in *N. benthamiana*. A link between ROS and HR during non-host resistance has been reported in lettuce-*Pseudomonas syringae* pv. *phaseolicola* (Bestwick *et al.* 1997) and tobacco-*Xanthomonas campestris* pv. *vesicatoria* pathosystems (Zurbriggen *et al.* 2009). Yoda *et al.* (2003) revealed that HR induction in tobacco plants inoculated with tobacco mosaic virus is through hydrogen peroxide produced by polyamine degradation. In conclusion, the results of this study demonstrated that HR induction in *N. tabacum* against *X. perforans* is affected by the inoculum concentration and the temperature of plant incubation post-inoculation. Additionally, faster and stronger HR elicitation by exogenous application of H₂O₂ confirmed the role of ROS in HR development. Reduction and delay in HR induction after exogenously supplied catalase also indicate the correlation between ROS accumulation and HR development.

References

- Alvarez M.E., Pennell R.I., Meijer P.J., Ishikawa A., Dixon R.A., Lamb C. 1998. Reactive oxygen intermediates mediate a systemic signal network in the establishment of plant immunity. *Cell* 92 (6): 773–784. DOI: 10.1016/S0092-8674(00)81405-1
- Bestwick C.S., Brown I.R., Bennett M., Mansfield J.W. 1997. Localization of hydrogen peroxide accumulation during the hypersensitive reaction of lettuce cells to *Pseudomonas syringae* pv. *phaseolicola*. *Plant Cell* 9 (2): 209–221. DOI: 10.1105/tpc.9.2.209
- Bolwell G.P., Daudi A. 2009. Reactive oxygen species in plant-pathogen interactions. p. 113–133. In: "Reactive Oxygen Species in Plant Signaling (Signaling and Communication in Plants)" (L.A. del Río, A. Puppo, eds.). Springer-Verlag, Berlin Heidelberg, Germany, 246 pp.
- Bradley D.J., Kjellbom P., Lamb C.J. 1992. Elicitor- and wound-induced oxidative cross-linking of a proline-rich plant cell wall protein: a novel, rapid defense response. *Cell* 70 (1): 21–30. DOI: 10.1016/0092-8674(92)90530-P
- Budde I.P., Ullrich M.S. 2000. Interactions of *Pseudomonas syringae* pv. *glycinea* with host and nonhost plants in relation to temperature and phytotoxin synthesis. *Molecular Plant-Microbe Interaction* 13 (9): 951–961. DOI: 10.1094/MPMI.2000.13.9.951
- Cheng Y., Zhang H., Yao J., Wang X., Xu J., Han Q., Wei G., Huang L., Kang Z. 2012. Characterization of non-host resistance in broad bean to the wheat stripe rust pathogen. *BMC Plant Biology* 12 (1): 96. DOI: 10.1186/1471-2229-12-96.
- Fan J., Doerner P. 2012. Genetic and molecular basis of non-host disease resistance: complex, yes; silver bullet, no. *Current Opinion in Plant Biology* 15 (4): 400–406. DOI: 10.1016/j.pbi.2012.03.001
- Hao X., Yu K., Ma Q., Song X., Li H., Wang M. 2011. Histochemical studies on the accumulation of H₂O₂ and hypersensitive cell death in the non-host resistance of pepper against *Blumeria graminis* f. sp. *tritici*. *Physiological and Molecular Plant Pathology* 76 (2): 104–111. DOI: 10.1016/j.pmp.2011.07.003
- Hückelhoven R., Dechert C., Kogel K.H. 2001. Non-host resistance of barley is associated with a hydrogen peroxide burst at sites of attempted penetration by wheat powdery mildew fungus. *Molecular Plant Pathology* 2 (4): 199–205. DOI: 10.1046/j.1464-6722.2001.00067.x.
- Jahnen W., Hahlbrock K. 1988. Cellular localization of nonhost resistance reactions of parsley (*Petroselinum crispum*) to fungal infection. *Planta* 173 (2): 197–204. DOI: 10.1007/BF00403011.
- Jones J.B., Lacy G.H., Bouzar H., Stall R.E., Schaad N.W. 2004. Reclassification of the xanthomonads associated with bacterial spot disease of tomato and pepper. *Systematic and Applied Microbiology* 27 (6): 755–762. DOI: 10.1078/0723202042369884
- Klement Z., Farkas G.L., Lovrekovich L. 1964. Hypersensitive reaction induced by phytopathogenic bacteria in the tobacco leaf. *Phytopathology* 54: 474–477.
- Kwak Y.S., Han K.S., Lee J.H., Lee K.H., Chung W.S., Mysore K.S., Kwon Y.S., Kim H.K., Bae D.W. 2009. Different oxidative burst pattern occur during host and non-host resistance responses triggered by *Xanthomonas campestris* in pepper. *Journal of Plant Biotechnology* 36: 244–254.
- Lamb C., Dixon R.A. 1997. The oxidative burst in plant disease resistance. *Annual Review of Plant Physiology* 48: 251–275. DOI: 10.1146/annurev.arplant.48.1.251
- Levine A., Tenhaken R., Dixon R., Lamb C. 1994. H₂O₂ from the oxidative burst orchestrates the plant hypersensitive disease resistance response. *Cell* 79 (4): 583–593. DOI: 10.1016/0092-8674(94)90544-4
- Li W., Xu Y.P., Yang J., Chen G.Y., Cai X.Z. 2015. Hydrogen peroxide is indispensable to *Xanthomonas oryzae* pv. *oryzae*-induced hypersensitive response and nonhost resistance in *Nicotiana benthamiana*. *Australasian Plant Pathology* 44: 611–617. DOI 10.1007/s13313-015-0376-1
- Mellersh D.G., Foulds I.V., Higgins V.J., Heath M.C. 2002. H₂O₂ plays different roles in determining penetration failure in

- three diverse plant-fungal interactions. *The Plant Journal* 29 (3): 257–268. DOI: 10.1046/j.0960-7412.2001.01215.x
- Mittler R., Vanderauwera S., Gollery M., Van Breusegem F. 2004. Reactive oxygen gene network of plants. *Trends in Plant Science* 9 (10): 490–498. DOI: 10.1016/j.tplants.2004.08.009
- Osdaghi E., Taghavi S.M., Hamzehzarghani H., Fazliarab A., Lamichhane J.R. 2016. Monitoring the occurrence of tomato bacterial spot and range of the causal agent *Xanthomonas perforans* in Iran. *Plant Pathology*. DOI: 10.1111/ppa.12642.
- Safaie Farahani A., Taghavi S.M. 2015. Expression profiling of malate dehydrogenase, superoxide dismutase and polygalacturonase-inhibiting protein in common bean in response to host and non-host pathogens. *Journal of Plant Pathology* 97 (3): 491–495. DOI: 10.4454/JPP.V97I3.030
- Safaie Farahani A., Taghavi M. 2016. Changes of antioxidant enzymes of mung bean [*Vigna radiata* (L.) R Wilczek] in response to host and non-host bacterial pathogens. *Journal of Plant Protection Research* 56 (1): 95–99. DOI: 10.1515/jppr-2016-0016
- Senthil-Kumar M., Mysore K.S. 2013. Non-host resistance against bacterial pathogens: retrospectives and prospects. *Annual Review of Phytopathology* 51: 407–427. DOI: 10.1146/annurev-phyto-082712-102319.
- Yoda H., Yamaguchi Y., San H. 2003. Induction of hypersensitive cell death by hydrogen peroxide produced through polyamine degradation in tobacco plants. *Plant Physiology* 132: 1973–1981. DOI: 10.1104/pp.103.024737
- Zurbriggen M.D., Carrillo N., Tognetti V.B., Melzer M., Peisker M., Hause B., Hajirezaei M.R. 2009. Chloroplast-generated reactive oxygen species play a major role in localized cell death during the non-host interaction between tobacco and *Xanthomonas campestris* pv. *vesicatoria*. *Plant Journal* 60 (6): 962–973. DOI: 10.1111/j.1365-313X.2009.04010.x.