

REVIEW

Interactions between entomopathogenic nematodes and entomopathogenic fungi in the aspect of new possibilities for biological plant protection

Katarzyna Barszczewska*  Oliwia Pietruszyńska

Department of Biological Methods, Institute of Plant Protection – National Research Institute, Poznań, Poland

Vol. 65, No. 3: 303–310, 2025

DOI: 10.24425/jppr.2025.155781

Received: September 27, 2024

Accepted: January 10, 2025

Online publication: September 08, 2025

*Corresponding address:

k.barszczewska@iorpib.poznan.pl

Responsible Editor:

Jolanta Kowalska

Abstract

Entomopathogenic fungi (EPF) and entomopathogenic nematodes (EPNs) have a strategic and important role in the biological control of agrophages. Micro- and macro-organisms have become an alternative to chemical methods, not only reducing the use of chemical pesticides, but also preventing agrophages from becoming resistant to chemicals and minimizing risks to human, animal and environmental health. The common habitat of EPNs and EPF is the soil, but interactions between them have mainly been studied in the laboratory. Recently, there has been a growing interest in combining biological control agents to increase their efficacy, and many studies have focused on combining EPF and EPNs against different insect pests. Studies have shown synergistic, additive or antagonistic effects. The results of these interactions depend on the pathogen, the host species, the different times of infection and the choice of the appropriate pathogen dose. The aim of this review was to summarize the existing knowledge on the life cycle of nematodes and the mechanisms of action of EPF and their application in practice, as well as the interaction between EPNs and EPF in plant protection against insect pests.

Keywords: biological control, entomopathogens, fungi, interactions, nematodes

Introduction

Intense environmental changes in recent years are posing a major challenge to modern agriculture. A warming climate, lower availability of fresh water, and emerging armed-political conflicts resulting in declining cultivated areas are becoming a problem. In addition, intensive agricultural production contributes to the degradation of arable land by reducing organic matter, nutrients, and biodiversity (Gullino *et al.* 2021). Large-scale farming makes it easier for diseases and pests to spread, and the high degree of chemicalization of agriculture is creating pesticide-resistant organisms and exacerbating food safety concerns (Premanandh 2011; Savary *et al.* 2019). Gaugler and Kaya (1990) showed that applied insecticides can enter the food chain and have long-term effects on organisms,

including humans. Since chemical pesticides pose a risk to human health and the environment, new options for crop protection need to be found. Alternatives to chemicals are biopesticides, which are based on the use of living organisms, i.e., micro-organisms (bacteria and fungi), viruses or macro-organisms (parasitic nematodes, predatory mites, and parasitic and predatory insects). These organisms can be introduced into the environment by an augmentative method, involving the introduction of a large number of beneficial micro- or macro-organisms that are naturally occurring enemies of pests (Collier and Van Steenwyk 2004). Organic food production is becoming increasingly popular in Europe, including Poland, and the area under organic crops is gradually increasing.

Biopreparations, including those containing EPNs or EPF, are gaining acceptance in plant protection (Kruk and Dziągiewlewska 2019; Sharapova 2019). Fungi and nematodes belong to a group of biotic agents that play several important roles in the environment and influence human life and activities. The biodiversity of microorganisms present influences the balance of individual environments. These include fungi and nematodes that parasitize other organisms and act as natural bioregulators in the environment. EPNs include the genera *Oscheius*, *Steinernema* and *Heterorhabditis* (Torrini *et al.* 2015; Poinar 2018). Their juveniles in the infective juvenile larva (IJs) stage live in the soil and are responsible for actively seeking out the host insect. Furthermore, IJs can survive at temperatures between 5°C and 35°C without losing their pathogenicity. EPNs are associated with bacteria, that once released into the insect's body, cause death within 24-48 hours (Gaugler 2002). Equally important bio-regulators of plant pests are EPF, which infect insects mainly in the soil environment. The most common EPF in the environment are *Entomophthora*, while biopesticides belonging to the genera *Beauveria*, *Cordyceps* (*Isaria*) and *Metarhizium* are most frequently used. Due to the wide range of infected hosts, EPF are found in aquatic and soil environments in all climatic zones. The wide range of occurrence has led to the development of a significant biodiversity (Araújo and Hughes 2016). The process of infecting prey involves the fungal spore entering the insect's body surface, germinating, and growing through the epidermis or chitinous exoskeleton. Under conditions of optimal humidity and temperature, the fungal hyphae completely colonize the insect's body. Depending on their morphology, the fungi form yeast-like threads, protoplasts, or filaments inside the host (Araújo and Hughes 2016). Most EPF kill their host before they can produce spores. The entire infection process takes about 4–7 days (Litwin *et al.* 2020; Quesada-Moraga *et al.* 2020). The combined use of EPF fungi and EPNs could be an innovative approach to pest control. Research to date has mainly been based on laboratory studies and have primarily focused on the insecticidal efficacy of fungi and nematodes, while their interactions and possible effects under natural conditions are largely unknown. Most research has reported an additive effect of the combination of EPNs and EPF on the mortality of some insect species, but in some cases an antagonistic interaction was found (Barbercheck and Kaya 1991; Ansari *et al.* 2004; Shapiro-Ilan *et al.* 2004; Tarasco *et al.* 2011; Půža and Tarasco 2023). Due to the different defense mechanisms of the pest, it is necessary to try to combine the use of EPF and EPNs. The entomopathogenic activity of these organisms is based on different actions, so the likelihood

of an insect developing defense mechanisms against several pathogenicity mechanisms at the same time is very low (Spescha *et al.* 2023). The development of EPF and EPNs resistance is therefore unlikely to pose an environmental risk in the future.

Mechanisms of action of entomopathogenic fungi and their use in practice

Entomopathogenic fungi are a special group of soil-dwelling microorganisms that infects and kills insects through cuticle penetration (Pell *et al.* 2010). Fungi have an important role as reductants, contributing to the decomposition of organic compounds and promoting the cycling of elements in the environment (Gadd 2007). A significant proportion of fungi cause plant diseases. Pathogens can weaken and damage crops, causing economic losses in agriculture and food production (Thambugala *et al.* 2020). The polyphagous nature of some fungal species affects their rapid spread and makes effective control difficult. Environmentally relevant parasitic fungal species are used for biological control (Singkaravanit *et al.* 2010). EPF are found in the Ascomycota, Zygomycota, Deuteromycota, Chytridiomycota and Oomycota clusters. The life cycle varies from group to group but is primarily based on infecting the host and causing death (Kidanu and Hagos 2020). All EPF are transmitted by spores (Araújo and Hughes 2016). The infection process begins with the attachment of the spores to the insect's epidermis, facilitated by the appressorium. This is the filamentous structure responsible for attaching the germinating spore to the surface of the host's epidermis. Penetration of the epidermis and growth of fungal hyphae inside the prey's body follows, along with secretion of lipases, proteins and chitinases. Insects may respond to the presence of entomopathogenic fungi in their bodies through several physiological mechanisms, e.g., immune and antioxidant responses (Aghaeepour *et al.* 2023). The host dies as a result of mechanical damage caused by the proliferating hyphae and by poisoning with toxins produced by the pathogen (Kidanu and Hagos 2020). Secondary metabolites produced during infection, including destruxins and cytochalasins, may be directly involved in killing the host or suppressing its defense responses, thereby indirectly aiding the pathogen (Fig. 1) (Vilcinskis *et al.* 1997).

A common phenomenon among fungi is antagonism – the substances they produce reduce the growth of other microorganisms, including fungi. There are known cases of antagonistic effects of entomopathogenic fungi on pathogenic fungi such as *Rhizoctonia*




Fig. 1. Life cycle of entomopathogenic fungi: 1 – infective stage – spore, 2 – spore adhesion to the epidermis, 3 – spore germination, 4 – growth of fungal hyphae inside the host body, 5 – death of the host and appearance of the outside mycelium with spores

solani Kühn and *Pythium myriotylum* Drechsler (Ownley *et al.* 2008). Some mycotoxins may have nematocidal activity. Beauvericin, secreted by fungi of the species *Beauveria bassiana* (Bals. – Criv.) Vuill. and *Beauveria pseudobassiana* (S.A. Rehner and Humber), has been shown to significantly reduce the incidence of the pine wood nematode (*Bursaphelenchus xylophilus* Steiner and Buhner), with up to 100% nematocidal efficacy in some variants (Sánchez-Gómez *et al.* 2023). Insect-pathogenic fungi play an important role in reducing plant pests. In the environment, aphid epizootics caused by fungi of the order Entomophthorales are frequently observed (Bałazy *et al.* 2008). Unfortunately, this species has no use in biopesticides. In practice, EPF are mainly used to control pests in cover crops, where the conditions are most favorable for their development (Ekbom 1979). For example, high pathogenicity of *B. bassiana* was observed against mealworms (*Tenebrio molitor* L.) and brown marmorated sting bug (*Halyomorpha halys* Stål). Within 10 days of application, 100% insect mortality was achieved under laboratory conditions (Swathy *et al.* 2024). Dannon *et al.* (2020) in their laboratory and greenhouse studies showed a significant effect of *B. bassiana* application on western flower thrips, *Frankliniella occidentalis* (Pergande), with a result of 69–96% insect mortality. Kuźniar *et al.* (2014) showed a positive effect of using *Isaria fumosorosea* (Wize) in faba bean (*Vicia faba* L.) cultivation, where the fungus significantly reduced the damage caused by the feeding of the pea leaf weevil (*Sitona lineatus* L.) on plant roots. In recent years, there has been a significant increase in the use of EPF in field crops, linked to the use of integrated pest management and an increasing range of biopreparations containing different strains of insect-pathogenic fungi (Kidanu and Hagos 2020).

The life cycle of entomopathogenic nematodes and their use in practice

For commercial pest control, nematodes belonging to the genera *Steinernema* and *Heterorhabditis* are used because of their wide host range, short developmental cycle, and ease of mass production *in vitro*. (Askary and Abd-Elgawad 2017). The life cycle of insect nematodes consists of the following stages: egg, four larval stages and adult. Species belonging to the genus *Steinernema* are dioecious, while in the genus *Heterorhabditis* the first generation is hermaphroditic and subsequent generations are dioecious. The stage that occurs in the soil is the infective juvenile larva (IJ) stage. This stage has the ability to actively seek out the host insect. These larvae enter the insect's body cavity through natural openings, i.e., mouth, anus or fistulae, and, in the case of species of the genus *Heterorhabditis*, by direct penetration of the insect's soft cuticle. After entering the insect host, they reach the haemocoel and release symbiotic bacteria from the gut. Nematodes of the *Steinernema* genus contain bacteria of the genus *Xenorhabdus*, while *Heterorhabditis* nematodes contain bacteria of the genus *Photorhabdus*. These bacteria release toxic and immunosuppressive compounds that kill the insect (Koppenhöfer and Gaugler 2009; Liao *et al.* 2017). When the insect dies, its body is bioconverted into a nutrient medium that helps the nematode grow and reproduce (Fig. 2). During this stage, the bacteria secrete small molecule antibiotics




Fig. 2. Life cycle of entomopathogenic nematodes of the genera *Steinernema* and *Heterorhabditis*

that protect the dead insect from other bacteria. Nematodes, together with symbiotic bacteria, are able to suppress the insect's immune system and weaken the antimicrobial response, e.g., lysozyme and cecropin are inhibited by *Xenorhabdus* species (Park *et al.* 2007; Lalitha *et al.* 2018). Characteristically, insects infected with nematodes of the genus *Heterorhabditis* turn a reddish maroon color. EPNs and their mutualistic bacteria are used commercially as a safe alternative to chemical insecticides. EPNs such as: *Steinernema carpocapsae* (Weiser), *Steinernema feltiae* (Filipjev), *Steinernema kraussei* (Steiner), *Steinernema glaseri* (Steiner), *Steinernema riobrave* (Cabanillas, Poinar and Raulston), *Heterorhabditis bacteriophora* (Poinar) and *Heterorhabditis megidis* (Poinar, Jackson and Klein) are among the species that are most often produced on an industrial scale. Through liquid culture, this has low production costs, for example, *H. bacteriophora* and *H. megidis* are successfully used against the black vine weevil (*Otiorhynchus sulcatus* Fabricius), which is considered a dangerous pest of yew, rhododendron, and many other nursery plants (Ehlers 2007). In the case of nematodes of the genus *Steinernema*, the genus *S. glaseri* for grubs, *S. kraussei* for the spruce web-spinning sawfly and *S. carpocapsae* for the *Hylobius* root weevils have been successfully recognized on the market (Shapiro-Ilan *et al.* 2020). EPNs are an effective and environmentally friendly pest control tool. Their versatility, ability to integrate with other crop protection methods (e.g., the use of pheromone traps, light traps or other biological control agents) and environmental benefits make them a valuable component of sustainable agriculture and crop protection programs (Oliveira-Hofman *et al.* 2019).

Interactions between entomopathogenic nematodes and entomopathogenic fungi

Biopesticides allow the combined use of a wide variety of pathogenic organisms. In recent years there has been increased interest in this mode of action, particularly in the combination of EPF strains with numerous species of EPNs against a wide range of insect pests. The effect of the combined use of entomopathogens can be antagonistic, where the presence of one pathogen inhibits the activity of other organisms, or synergistic, where one pathogen facilitates the infection of other organisms (Fig. 3) (Correa-Cuadros *et al.* 2016; Wakil *et al.* 2017). Most of the studies conducted on the combined use of EPF and EPN have been based on the control of beetles of the families Scarabaeidae and Curculionidae (Coleoptera) and insects of the order Lepidoptera (Shapiro-Ilan *et al.* 2004;




Fig. 3. Schematic of the interaction between entomopathogenic fungi and entomopathogenic nematodes

Otsuki and Yano 2014). One example is research into the combination of the fungi *B. bassiana* and *Metarhizium anisopliae* (Metchnikoff) with the nematode *H. bacteriophora* against the larvae of the palm weevil (*Rhynchophorus ferrugineus* Olivier). Additive and synergistic interactions were found in these studies. It was also found that a 1–2-week delay in nematode application after the fungus increased the efficacy of the treatment, as the longer the larvae are exposed to the fungus, the more weakened they become and consequently more susceptible to EPNs (Wakil *et al.* 2017). In the case of insects of the order Lepidoptera, a combination of *B. bassiana* and *H. bacteriophora* was used resulting in an additive effect, causing a higher mortality rate of beet armyworm larvae (*Spodoptera exigua* Hübner). To achieve this effect, a double infestation of nematodes and a fungus was carried out (Barbercheck and Kaya 1991). Ansari *et al.* (2006) observed in field experiments on white grubs (*Hoplia philanthus* Fuessly, Coleoptera: Scarabaeidae) that the combined application of *M. anisopliae* and *H. megidis* resulted in additive or synergistic effects. More than 95% of larval mortality was observed following nematode application 4 weeks after fungal application. Similar additive effects of *Heterorhabditis indica* (Poinar, Karunakar and David) and *M. anisopliae* were shown by Shapiro-Ilan *et al.* (2004) in experiments on pecan weevil larvae (*Curculio caryae* Grub, Coleoptera: Curculionidae). When the fungus *B. bassiana* was used together with *H. indica*, the interactions were antagonistic, as in the variant using *S. carpocapsae* and *Paecilomeces fumosoroseus* (*I. fumosorosea*) Wize.

Studies on EPF and EPNs show that the efficacy of entomophagy depends on a number of abiotic factors. It has been observed that the optimum temperature for spore germination of entomopathogenic fungi depends on the region of origin of the strain and its species, but 20–30°C is considered a favorable temperature range (Gul *et al.* 2014). An important element in the case of fungi is adequate ambient humidity. The most favorable conditions are those where the relative humidity is close to saturation (>95.5%) and this is the

range in which the highest level of spore germination is achieved (Abdul Qayyum *et al.* 2021). In the case of EPNs, a broader spectrum of action is shown. Some studies report that lower soil moisture can negatively affect nematode pathogenicity. Maintaining adequate

moisture conditions allows the nematodes to move more efficiently and increases their survival rate (Toth *et al.* 2022). The rate of growth and development of EPNs and EPF depends on the substrate in which they occur. Studies by Bueno-Pallero *et al.* (2018) show that

Table 1. Interactions between entomopathogenic fungi and entomopathogenic nematodes during application to agricultural pests (Půža and Tarasco 2023)

Target Pest	Nematode	Fungus	Interaction
<i>Bactrocera dorsalis</i> (Diptera: Tephritidae)	<i>H. bacteriophora</i> <i>S. carpocapsae</i>	<i>B. bassiana</i> <i>M. anisopliae</i>	additive, synergistic
<i>Bactrocera zonata</i> (Diptera: Tephritidae)	<i>H. bacteriophora</i> <i>S. carpocapsae</i>	<i>B. bassiana</i> <i>M. anisopliae</i>	additive, synergistic
<i>Coptognathus curtispennis</i> (Coleoptera: Scarabaeidae)	<i>H. bacteriophora</i> <i>S. yirgalemense</i>	<i>M. anisopliae</i>	additive, synergistic
<i>Curculio caryae</i> (Coleoptera: Curculionidae)	<i>H. indica</i> <i>S. carpocapsae</i>	<i>B. bassiana</i> <i>M. anisopliae</i> <i>C. fumosorosea</i>	antagonistic, additive
<i>Curculio nucum</i> (Coleoptera: Curculionidae)	<i>H. bacteriophora</i> <i>S. carpocapsae</i> <i>S. feltiae</i>	<i>M. anisopliae</i>	additive
<i>Cyclocephala lurida</i> (Coleoptera: Scarabaeidae)	<i>H. bacteriophora</i> <i>H. megidis</i>	<i>B. bassiana</i> <i>M. anisopliae</i>	additive, synergistic
<i>Diatraea saccharalis</i> (Lepidoptera: Pyralidae)	<i>H. bacteriophora</i>	<i>M. anisopliae</i>	additive, antagonistic
<i>Ectinohoplia rufipes</i> (Coleoptera: Scarabaeidae)	<i>H. bacteriophora</i> <i>S. carpocapsae</i>	<i>B. brongniartii</i>	additive
<i>Exomala orientalis</i> (Coleoptera: Scarabaeidae)	<i>H. bacteriophora</i> <i>S. carpocapsae</i>	<i>B. brongniartii</i>	additive
<i>Hoplia philanthus</i> (Coleoptera: Scarabaeidae)	<i>H. bacteriophora</i> <i>H. megidis</i> <i>S. glaseri</i>	<i>M. anisopliae</i>	additive, synergistic
<i>Hylobius abietis</i> (Coleoptera: Curculionidae)	<i>S. carpocapsae</i> <i>H. downesi</i>	<i>B. bassiana</i> <i>M. anisopliae</i> <i>M. brunneum</i> <i>B. caledonica</i>	additive
<i>Leptinotarsa decemlineata</i> (Coleoptera: Chrysomelidae)	<i>S. feltiae</i> <i>H. bacteriophora</i>	<i>B. bassiana</i> <i>C. fumosorosea</i>	additive
<i>Otiorhynchus sulcatus</i> (Coleoptera: Curculionidae)	<i>H. bacteriophora</i> <i>S. carpocapsae</i> <i>S. kraussei</i>	<i>M. anisopliae</i>	additive, synergistic
<i>Plutella xylostella</i> (Lepidoptera: Plutellidae)	<i>H. bacteriophora</i>	<i>B. bassiana</i> <i>M. anisopliae</i>	antagonistic, additive, synergistic
<i>Rhagoletis pomonella</i> (Diptera: Tephritidae)	<i>S. carpocapsae</i> <i>S. riobrave</i>	<i>B. bassiana</i> <i>M. brunneum</i> <i>C. javanica</i> <i>C. fumosorosea</i>	additive
<i>Rhynchophorus ferrugineus</i> (Coleoptera: Curculionidae)	<i>H. bacteriophora</i>	<i>B. bassiana</i> <i>M. anisopliae</i>	additive, synergistic
<i>Spodoptera exigua</i> (Lepidoptera: Noctuidae)	<i>H. bacteriophora</i> <i>S. carpocapsae</i>	<i>B. bassiana</i>	additive
<i>Spodoptera littoralis</i> (Lepidoptera: Noctuidae)	<i>H. bacteriophora</i> <i>S. riobrave</i>	<i>B. bassiana</i>	synergistic
<i>Thaumetotibia leucotreta</i> (Lepidoptera: Tortricidae)	<i>S. jeffreyense</i> <i>S. yirgalemense</i>	<i>M. pinghaense</i>	antagonistic, additive, synergistic
<i>Thrips tabaci</i> (Thysanoptera: Thripidae)	<i>H. bacteriophora</i> <i>S. feltiae</i>	<i>B. bassiana</i> <i>M. anisopliae</i>	additive

the growth effect on a limiting substrate (CMA ¼) is slower than on a rich substrate (PDA), indicating that the availability of resources in the soil plays an important role in the growth and development of EPNs and the growth of EPF. Several biotic and abiotic factors also determine the epizootic capacity of EPF (Mudgal *et al.* 2013). Among the biotic factors, the nematode strain, the density of invasive intestinal nematodes, nematode strain, density of invading IJs, insect host species and insect host age and nutritional status of the host insects have a significant impact on the interaction between EPNs and EPF, as they are linked to the persistence and development of nematode populations and thus influence the infectivity of EPNs (Shaurub 2023).

Secondary metabolites from *Xenorhabdus* and *Photorhabdus* (these bacteria are found in the gastrointestinal tract of nematodes of the genera *Steinernema* and *Heterorhabditis*) have been shown to be effective in competition between EPF and EPNs. For example, Tarasco *et al.* (2011) showed that secondary metabolites produced by *X. bovienii* inhibited the growth of *B. bassiana*. This is one of the main problems with using fungi and nematodes at the same time. Once inside the insect, EPN and EPF can release specific metabolites to protect the insect from other macro-organisms. With such strong competition in the insect haemocoel, exclusion of one of the organisms may occur (Strasser *et al.* 2000). In contrast, Ansari *et al.* (2008) used a combination of EPF *M. anisopliae* together with EPN – *H. bacteriophora*, *S. feltiae* and *S. kraussei*, which resulted in increased efficacy against black vine weevil (*Otiorhynchus sulcatus*). The application of EPN 1 or 2 weeks after fungus application resulted in 100% larval mortality, even when biological control agents were applied at reduced doses. Another positive effect of the combined application of EPF and EPN was shown by Usman *et al.* (2020). In experiments with *Rhagoletis pomonella* (Walsh, Diptera: Tephritidae), the combination of nematodes and fungi was shown to be more effective than either fungi or nematodes alone.

Current studies on the combined use of EPNs and EPF include seven fungal and 10 nematode species. The studies clearly show a significant effect of bioagents on the efficacy of their combined use (Table 1). Some studies have reported an antagonistic effect of pathogen combinations on application efficacy, but some combinations may have a negative effect on the propagation of both pathogens. Therefore, this aspect should play a key role when considering specific nematode-fungus combinations (Půža and Tarasco 2023).

References

- Abdul Qayyum M., Bilal H., Naeem Ullah U., Ali H., Raza H., Wajid M. 2021. Factors affecting the epizootics of entomopathogenic fungi – a review. *Journal of Bioresource Management* 8 (4): 78–85. DOI: 10.35691/JBM.1202.0204
- Aghaeepour S., Zibae A., Ramzi S., Hoda H. 2023. Changes in life table parameters and intermediary metabolism of *Cryptolaemus montrouzieri* Mulsant after infection by *Beauveria bassiana*. *Journal of Plant Protection Research* 63 (1): 68–82. DOI: 10.24425/jppr.2023.144503
- Ansari M.A., Shah F.A., Tirry L., Moens M. 2006. Field trials against *Hoplia philanthus* (Coleoptera: Scarabaeidae) with a combination of an entomopathogenic nematode and the fungus *Metarhizium anisopliae* CLO 53. *Biological Control* 39 (3): 453–459. DOI: 10.1016/j.biocontrol.2006.07.004
- Ansari M.A., Tirry L., Moens M. 2004. Interaction between *Metarhizium anisopliae* CLO 53 and entomopathogenic nematodes for the control of *Hoplia philanthus*. *Biological Control* 31 (2): 172–180. DOI: 10.1016/j.biocontrol.2004.04.002
- Ansari M.A., Shah F.A., Butt T.M. 2008. Combined use of entomopathogenic nematodes and *Metarhizium anisopliae* as a new approach for black vine weevil, *Otiorhynchus sulcatus*, control. *Entomologia Experimentalis et Applicata* 129 (3): 340–347. DOI: 10.1111/j.1570-7458.2008.00783.x
- Araújo J.P.M., Hughes D.P. 2016. Chapter one – diversity of entomopathogenic fungi: which groups conquered the insect body? *Advances in Genetics* 94: 1–39. DOI: 10.1016/bs.adgen.2016.01.001
- Askary T.H., Ahmad M.J. 2017. Entomopathogenic nematodes: mass production, formulation, and application. p. 261–286. In: “Biocontrol agents: entomopathogenic and slug parasitic nematodes” (Abd-Elgawad M.M.M., Askary T.H., Couland J., eds.). CABI, UK.
- Bałaży S., Miętkiewski R., Tkaczuk C., Wegensteiner R., Wrzosek M. 2008. Diversity of acaropathogenic fungi in Poland and other European countries. *Experimental and Applied Acarology* 46 (1): 53–70.
- Barbercheck M.E., Kaya H.K. 1991. Competitive interactions between entomopathogenic nematodes and *Beauveria bassiana* (Deuteromycotina: Hyphomycetes) in soilborne larvae of *Spodoptera exigua* (Lepidoptera: Noctuidae). *Environmental Entomology* 20 (2): 707–712. DOI: 10.1093/ee/20.2.707
- Bueno-Pallero F. Á., Blanco-Pérez R., Dionisio L., Campos-Herrera R. 2018. Simultaneous exposure of nematophagous fungi, entomopathogenic nematodes and entomopathogenic fungi can modulate belowground insect pest control. *Journal of Invertebrate Pathology* 154: 85–94. DOI: 10.1016/j.jip.2018.04.004
- Collier T., Van Steenwyk R. 2004. A critical evaluation of augmentative biological control. *Biological Control* 31 (2): 245–256. DOI: 10.1016/j.biocontrol.2004.05.001
- Correa-Cuadros J.P., Sáenz-Aponte A., Rodríguez-Bocanegra M.X. 2016. *In vitro* interaction of *Metarhizium anisopliae* Ma9236 and *Beauveria bassiana* Bb9205 with *Heterorhabditis bacteriophora* HNI0100 for the control of *Plutella xylostella*. *SpringerPlus* 5: 1–8. DOI: 10.1186/s40064-016-3745-5
- Dannon H.F., Dannon A.E., Douro-Kpindou O.K., Zinsou A.V., Houndete A.T., Toffa-Mehinto J., Elegbede I.A.T.M., Olou B.D., Tamò M. 2020. Toward the efficient use of *Beauveria bassiana* in integrated cotton insect pest management. *Journal of Cotton Research* 3: 1–21. DOI: 10.1186/s42397-020-00061-5
- Ehlers R.U. 2007. Nematodes: from Science to Commercial Use. p. 135–151. In: “Biological control: a global perspective: case studies from around the world” (C. Vincent, M.S. Goeettel, G. Lazarovits, eds.). CABI International. Kiel, Germany, 24118 pp.
- Ekbom B.S. 1979. Investigations on the potential of a parasitic fungus *Verticillium lecanii* for biological control of the greenhouse whitefly (*Trialeurodes vaporariorum*). *Swedish Journal of Agriculture Research* 9: 129–138.
- Gadd G.M. 2007. Geomycology: biogeochemical transformations of rocks, minerals, metals, and radionuclides by fungi, bioweathering and bioremediation. *Mycological Research*

- 111 (1): 3–49. DOI: 10.1016/j.mycres.2006.12.001
- Gaugler R., Kaya H.K. 1990. Entomopathogenic Nematodes in Biological Control. Boca Raton, CRC Press, FL, USA.
- Gaugler R. 2002. Entomopathogenic Nematology. CABI Publishing: Wallingford, UK, 388 pp.
- Gul H.T., Saeed S., Khan F.Z.A. 2014. Entomopathogenic fungi as effective insect pest management tactic: a review. Applied Sciences and Business Economics 1 (1): 10–18.
- Gullino M.L., Albajes R., Al-Jboory I., Angelotti F., Chakraborty S., Garrett K.A., Hurley B. P., Juroszek P., Makkouk K., Pan X., Stephenson T. 2021. Scientific review of the impact of climate change on plant pests – A global challenge to prevent and mitigate plant pest risks in agriculture, forestry, and ecosystems. IPPC Secretariat. Rome, Italy. DOI: 10.4060/cb4769en
- Kidanu S., Hagos L. 2020. Research and application of entomopathogenic fungi as pest management option: a review. Journal of Environment and Earth Science 10 (3): 31–39. DOI: 10.7176/JEES/10-3-03
- Koppenhöfer H.S., Gaugler R. 2009. Entomopathogenic nematode and bacteria mutualism. p. 99–116. In: “Defensive Mutualism in Microbial Symbiosis” (J. White, M. Torres, eds.). CRC Press, Boca Raton, Florida, USA.
- Kruk K., Dzięgielewska M. 2019. Wykorzystanie nicieni owadobójczych w biologicznej ochronie roślin. [The use of entomopathogenic nematodes in biological plant protection]. Młodzi Naukowcy 11: 72–76.
- Kuźniar T., Ropek D., Kulig B. 2014. Wykorzystanie grzyba owadobójczego *Isaria fumosorosea* do zwalczania szkodników w uprawie bobiku. Proceedings of ECOpole 8 (1): 201–207. DOI: 10.2429/proc.2014.8(1)026
- Lalitha K., Karthi S., Vengateswari G., Karthikraja R., Perumal P., Shivakumar M.S. 2018. Effect of entomopathogenic nematode of *Heterorhabditis indica* infection on immune and antioxidant system in lepidopteran pest *Spodoptera litura* (Lepidoptera: Noctuidae). Journal of Parasitic Diseases 42 (2): 204–211. DOI: 10.1007/s12639-018-0983-1
- Liao C., Gao A., Li B., Wang M., Shan L. 2017. Two symbiotic bacteria of the entomopathogenic nematode *Heterorhabditis* spp. against *Galleria mellonella*. Toxicon 127: 85–89. DOI: 10.1016/j.toxicon.2016.11.257
- Litwin A., Nowak M., Różalska S. 2020. Entomopathogenic fungi: unconventional applications. Reviews in Environmental Science and Biotechnology 19: 23–42. DOI: 10.1007/s11157-020-09525-1
- Mudgal S., Toni A., Tostivint C., Hokkanen H., Chandler D. 2013. Scientific support, literature review and data collection and analysis for risk assessment on microbial organisms used as active substance in plant protection products – Lot 1 Environmental Risk characterization. EFSA Supporting Publication 2013, 149 pp. DOI: 10.2903/sp.efsa.2013.EN-518.
- Oliveira-Hofman C., Kaplan F., Stevens G., Lewis E., Wu S., Alborn H.T., Shapiro-Ilan D.I. 2019. Pheromone extracts act as boosters for entomopathogenic nematodes efficacy. Journal of Invertebrate Pathology 164: 38–42. DOI: 10.1016/j.jip.2019.04.008
- Otsuki H., Yano S. 2014. Functionally different predators break down antipredator defenses of spider mites. Entomologia Experimentalis et Applicata 151 (1): 27–33. DOI: 10.1111/eea.12164
- Ownley B.H., Griffin M.R., Klingeman W.E., Gwinn K.D., Moulton J.K., Pereira R.M. 2008. *Beauveria bassiana*: endophytic colonization and plant disease control. Journal of Invertebrate Pathology 98: 267–270. DOI: 10.1016/j.jip.2008.01.010
- Park Y., Herbert E.E., Cowles C.E., Cowles K.N., Menard M.L., Orchard S.S., Goodrich-Blair H. 2007. Clonal variation in *Xenorhabdus nematophila* virulence and suppression of *Manduca sexta* immunity. Cellular Microbiology 9 (3): 645–656. DOI: 10.1111/j.1462-5822.2006.00815.x
- Pell J.K., Hannam J.J., Steinkraus D.C. 2010. Conservation biological control using fungal entomopathogens. BioControl 55 (1): 187–198. DOI: 10.1007/s10526-009-9245-6
- Poinar G.O. 2018. Nematodes for Biological Control of Insects. CRC Press: Boca Raton, FL, USA, 289 pp.
- Premanandh J. 2011. Factors affecting food security and contribution of modern technologies in food sustainability. Journal of the Science of Food and Agriculture 91 (15): 2707–2714. DOI: 10.1002/jsfa.4666
- Půža V., Tarasco E. 2023. Interactions between entomopathogenic fungi and entomopathogenic nematodes. Microorganisms 11 (1): 163. DOI: 10.3390/microorganisms11010163
- Quesada-Moraga E., Yousef-Naef M., Garrido-Jurado I. Advances in the use of entomopathogenic fungi as biopesticides in suppressing crop pests. p. 63–98. In: “Biopesticides for Sustainable Agriculture” (N. Brich, T. Glare, eds.). Burleigh Dodds Science Publishing: Cambridge, UK.
- Sánchez-Gómez T., Harte S.J., Zamora P., Bareyre M., Díez J.J., Herrero B., Niño-Sánchez J., Martín-García J. 2023. Nematicidal effect of *Beauveria* species and the mycotoxin beauvericin against pinewood nematode *Bursaphelenchus xylophilus*. Frontiers in Forests and Global Change 6: 1229456. DOI: 10.3389/ffgc.2023.1229456
- Savary S., Willocquet L., Pethybridge S.J., Esker P., McRobert N., Nelson A. 2019. The global burden of pathogens and pests on major food crops. Nature Ecology and Evolution 3: 430–439. DOI: 10.1038/s41559-018-0793-y
- Shapiro-Ilan D., Hazir S., Glazer I. 2020. Advances in use of entomopathogenic nematodes in integrated pest management. In: “Integrated management of insect pests: current and future developments” (M. Kogan, E.A. Heinrichs, eds.). Burleigh Dodds Science Publication, UK, 30 pp.
- Shapiro-Ilan D.I., Jackson M., Reilly C.C., Hotchkiss M.W. 2004. Effects of combining an entomopathogenic fungi or bacterium with entomopathogenic nematodes on mortality of *Curculio caryae* (Coleoptera: Curculionidae). Biological Control 30 (1): 119–126. DOI: 10.1016/j.biocontrol.2003.09.014
- Sharapova I.E. 2019. Prospects of using entomopathogenic fungus in development of a biopesticide product with nematocidal activity. Biocatalysis and Agricultural Biotechnology 19: 101098. DOI: 10.1016/j.bcab.2019.101098
- Shaurub E.S.H. 2023. Review of entomopathogenic fungi and nematodes as biological control agents of tephritid fruit flies: current status and a future vision. Entomologia Experimentalis et Applicata 171 (1): 17–34. DOI: 10.1111/eea.13244
- Singkaravanit S., Kinoshita H., Ihara F., Nihira T. 2010. Cloning and functional analysis of the second geranyl diphosphate synthase gene influencing helvolic acid biosynthesis in *Metarhizium anisopliae*. Applied Microbiology and Biotechnology 87 (3): 1077–1088. DOI: 10.1007/s00253-010-2556-9
- Spescha A., Weibel J., Wyser L., Brunner M., Hermida M.H., Moix A., Scheibler F., Guyer A., Campos-Herrera R., Grabenweger G., Maurhofer M. 2023. Combining entomopathogenic *Pseudomonas* bacteria, nematodes and fungi for biological control of a below-ground insect pest. Agriculture, Ecosystems and Environment 348: 108414. DOI: 10.1016/j.agee.2023.108414
- Strasser H., Abendstein D., Stuppner H., Butt T.M. 2000. Monitoring the distribution of secondary metabolites produced by the entomogenous fungus *Beauveria brongniartii* with particular reference to oosporein. Mycological Research 104 (10): 1227–1233. DOI: 10.1017/S0953756200002963
- Swathy K., Parmar M.K., Vivekanandhan P. 2024. Biocontrol efficacy of entomopathogenic fungi *Beauveria bassiana* conidia against agricultural insect pests. Environmental Quality Management 34 (1): e22174. DOI: 10.1002/tqem.22174

- Tarasco E., Santiago Alvarez C., Triggiani O., Quesada Moraga E. 2011. Laboratory studies on the competition for insect haemocoel between *Beauveria bassiana* and *Steinernema ichnusa* recovered in the same ecological niche. *Bio-control Science and Technology* 21 (6): 693–704. DOI: 10.1080/09583157.2011.570428
- Thambugala K.M., Daranagama D.A., Philips A.J.L., Kannangara S.D., Promputtha I. 2020. Fungi vs. Fungi in biocontrol: an overview of fungal antagonists applied against fungal plant pathogens. *Frontiers in Cellular and Infection Microbiology* 10: 604923. DOI: 10.3389/fcimb.2020.604923
- Torrini G., Mazza G., Carletti B., Benvenuti C., Roversi P.F., Fanelli E., De Luca F., Troccoli A. and Tarasco E. 2015. *Oscheius onirici* sp. n. (Nematoda: Rhabditidae): a new entomopathogenic nematode from an Italian cave. *Zootaxa* 3937 (3): 533–548. DOI:10.11646/zootaxa.3937.3.6
- Toth S., Toepfer S., Szalai M., Kiss J. 2022. Limited influence of abiotic and biotic factors on the efficacy of soil insecticides and entomopathogenic nematodes when managing the maize pest *Diabrotica v. virgifera* (Coleoptera: Chrysomelidae). *Agronomy* 12 (11): 2697. DOI: 10.3390/agronomy12112697
- Usman M., Gulzar S., Wakil S., Wu S., Piñero J.C., Leskey T.C., Nixon L.J., Oliveira-Hofman C., Toews M.D., Shapiro-Ilan D. 2020. Virulence of entomopathogenic fungi to *Rhagoletis pomonella* (Diptera: Tephritidae) and interactions with entomopathogenic nematodes. *Journal of Economic Entomology* 113 (6): 2627–2633. DOI: 10.1093/jee/toaa209
- Vilcinskis A., Matha V., Götz P. 1997. Effects of the entomopathogenic fungus *Metarhizium anisopliae* and its secondary metabolites on morphology and cytoskeleton of plasmacytes isolated from the greater wax moth, *Galleria mellonella*. *Journal of Insect Physiology* 43 (12): 1149–1159. DOI: 10.1016/S0022-1910(97)00066-8
- Wakil W., Yasin M., Shapiro-Ilan D. 2017. Effects of single and combined applications of entomopathogenic fungi and nematodes against *Rhynchophorus ferrugineus* (Olivier). *Scientific Reports* 7 (1): 5971. DOI: 10.1038/s41598-017-05615-3