

# DYNAMICS OF PREDATORY SYRPHIDAE IN THE APPLE ORCHARD AND NEIGHBOURING SHRUBBERIES

Paweł Trzciniński\*, Hanna Piekarska-Boniecka

Department of Entomology and Environmental Protection, Poznan University of Life Sciences  
Dąbrowskiego 159, 60-594 Poznań, Poland

Received: September 13, 2012

Accepted: March 15, 2013

**Abstract:** Shrubberies which neighbour on an orchard are among the significant structures enhancing agrocenoses. Such shrubberies maintain species diversity and stimulate self-control mechanisms in biocenoses. For these reasons, from 1999 to 2001, studies were conducted concerning predatory Syrphidae (Diptera) occurring in the orchard and shrubberies in the immediate vicinity. A quantity and quality analysis of predatory hoverflies (Diptera, Syrphidae) was performed. A total of 801 specimens were caught in both habitats. There were 20 species recorded there (12% of the national fauna of this predatory family).

The studies proved that the orchard and the neighbouring shrubberies can be alternative habitats for Syrphidae (Diptera). There is also a possibility of the predators migrating from the shrubberies to the orchard, enhancing the potential to control the abundant Hemiptera species in the orchard. The shrubberies then become an important element of the agrocenosis structure, due to their biocenotic functions.

**Key words:** orchard, predator, shrubberies, Syrphidae, Wielkopolska

## INTRODUCTION

Shrubberies surrounding orchards are vital structures enhancing agrocenoses. They provide shelter, a place for wintering and breeding as well as dispersion of entomophages. They are the feeding place and habitat for prey. They maintain species diversity and stimulate self-control mechanisms in agrocenoses (Bianchi *et al.* 2006; Debras *et al.* 2006; Haenke *et al.* 2009). A positive impact of the shrubberies and wild greenery of orchard borders on predatory Syrphidae (Diptera) in the orchard, was presented in studies by Solomon *et al.* (2000), Bostanian *et al.* (2004), Debras *et al.* (2006) and Rossi *et al.* (2006). Some papers dealing with this topic have been published in Polish, however, they presented the impact of wild plants of cultivated fields on Syrphidae occurrence (Galecka 1995; Kaczorowski and Dębek-Jankowska 1997; Garbarkiewicz and Trojanowski 1998; Bennewicz 2011). The preliminary results of research defining the influence of shrubberies which neighbour orchards, and the influence of the orchard itself on predators in those habitats, was presented in a paper by Trzciniński and Piekarska (2009).

The first aim of our study was to describe the changes in the quantity and quality structure of the predatory Syrphidae (Diptera) species occurring in the orchard and shrubberies in its vicinity. The second aim was to compare the communities of predators in the studied habitats in terms of the quantity and quality.

## MATERIALS AND METHODS

The study was carried out in the 1999–2001 time period, in the orchard of Przybroda in Poland (52°31'14"N, 16°39'22"E). Predators were caught from May to September in Moericke traps (Moericke 1953). Twelve traps were laid out in selected areas in the orchard and shrubberies and then checked every 10 days. The insects caught in 10 days, in one trap, constituted one sample.

The study area was divided into plots totalling 1.5 ha, where trees of the following fruit cultivars were grown: (a) apples: Primula, Novamac, Priam, Cortland, McIntosh; (b) sweet cherries: Red Büttner, Hedelwińska, and (c) plums: Anna Späth, Cacańska Rodna and Węgierka Dąbrowicka. The shrubberies adjacent to the orchard, formed a belt which was about 1 km long and 7–8 m wide. They were mainly: blackthorn plum-tree (*Prunus spinosa* L.), common pear-tree (*Pyrus communis* L.), common elm (*Ulmus campestris* L.), European spindle tree (*Evonymus europaea* L.), hawthorn (*Crataegus monogyna* Jacq.), whitethorn (*C. oxyacantha* L.), elder (*Sambucus nigra* L.), dewberry (*Rubus caesius* L.), blackberry (*R. suberectus* Anders.), briarrose (*Rosa canina* L.), common carrot (*Daucus carota* L.), common yarrow (*Achillea millefolium* L.), common nettle (*Urtica dioica* L.) and couch-grass (*Agropyron repens* L.).

The orchard followed an integrated fruit production policy, with 5–8 disease-prevention procedures and 6–8 anti-pest procedures performed in various study years.

The Syrphidae communities were characterised on the basis of the following parameters: dominancy index,

\*Corresponding address:  
trzcinsk@up.poznan.pl

coefficient of the general species diversity – H' (Shannon and Weaver 1963), and uniformity index J' (Pielou 1966). The Syrphidae communities were compared in quantities with the Re index (Renkonen 1938) and in terms of quality with the QS index (Sørensen 1948)

## RESULTS

In the 1999–2001 time period, 650 samples were collected from the orchard and 631 from the shrubberies. A total of 801 Syrphidae were caught in both habitats. There were more predators captured in the orchard. A total of 474 individuals were caught in the orchard and 327 in the shrubberies. In particular study years, similar fluctuations of the abundance of Syrphidae caught, were reported in both habitats. In 1999, the highest number re-

ported was 273 from the orchard and 200 from the shrubberies. The lowest number were caught in 2000, with 62 and 24 specimens, respectively (Table 1).

In both habitats, 20 Syrphidae were reported in total, *i.e.* 12% of the national fauna of this predatory family.

A similar species diversity in orchard and shrubberies was found, as 12 species were caught in the orchard and 14 in the shrubberies. Lower fluctuations in the orchard than in shrubberies were recorded in the number of species in particular years. In 1999 and 2001, the numbers of species were 10 and 7. In 2000 there were only 5 species. In the shrubberies, the fluctuations in the number of species caught were higher than in the orchard. In 1999, the number of species was 13, while only 4 species were caught in 2000, and also only 4 in 2001. (Table 1).

Table 1. The list of and abundance of Syrphidae species caught in the orchard and shrubberies, in Przybroda, 1999–2001

Species	Environment							
	orchard			Total	shrubberies			Total
	1999	2000	2001		1999	2000	2001	
<i>Baccha elongata</i> (Fabrucijs, 1775)	–	–	–	–	1	1	–	2
<i>Dasysyrphus venustus</i> (Meigen, 1822)	–	1	–	1	–	–	–	–
<i>Epistrophe eligans</i> (Harris, 1780)	1	–	–	1	–	–	–	–
<i>E. nitidicollis</i> (Meigen, 1822)	–	–	–	–	1	–	–	1
<i>Episyrphus balteatus</i> (De Geer, 1776)	258	58	105	421	175	21	100	296
<i>Eupeodes corollae</i> (Fabricius, 1794)	–	1	14	15	–	1	1	2
<i>Leucozona lucorum</i> (Linnaeus, 1758)	–	–	–	–	2	–	–	2
<i>Melanostoma mellinum</i> (Linnaeus, 1758)	2	–	3	5	1	–	1	2
<i>M. scalare</i> (Fabrucijs, 1794)	–	–	2	2	10	–	–	10
<i>Meliscaeva auricollis</i> (Meigen, 1822)	–	–	1	1	–	–	–	–
<i>Platycheirus albimanus</i> (Fabrucijs, 1781)	1	–	1	2	1	–	–	1
<i>P. peltatus</i> (Meigen, 1822)	–	–	–	–	2	–	–	2
<i>P. scutatus</i> (Meigen, 1822)	–	–	–	–	1	–	1	2
<i>Scaeva pyrastris</i> (Linnaeus, 1758)	2	1	1	4	1	–	–	1
<i>Sphaerophoria scripta</i> (Linnaeus, 1758)	–	–	1	1	–	–	–	–
<i>S. taeniata</i> (Meigen, 1822)	–	–	–	–	3	–	–	3
<i>Syrphus ribesii</i> (Linnaeus, 1758)	3	–	3	6	–	–	–	–
<i>S. torvus</i> Osten-Sacken, 1875	–	–	–	–	1	–	–	1
<i>S. vitripennis</i> Meigen, 1822	6	1	8	15	–	–	–	–
<i>Xanthandrus comtus</i> (Harris, 1780)	–	–	–	–	1	1	–	2
Total of specimens	273	62	139	474	200	24	103	327
	801							
Total of species	7	5	10	12	13	4	4	14

Among the predatory Syrphidae, only one species *Episyrphus balteatus* (De Geer), was the definite dominant. It occurred in large numbers in both habitats, with 88.8% in the group of orchard predators, and 90.5% in the shrubberies (Fig. 1, 2). The domination structure of the predatory Syrphidae caught in the orchard besides the eudominant *E. balteatus*, was also made up of 2 subdominant species, i.e. *E. corollae* (3.2%) and *Syrphus vitripennis* (3.2%), and the recedent *S. ribesii* (1.3%). A simpler domination structure of predatory Syrphidae was found in the shrubberies, because apart from the eudominant *E. balteatus* only one subdominant was caught, namely *M. scalare* (3%). Other species occurred sporadically and were only caught as single specimens both in the orchard and in the shrubberies.

Such a quantity-quality structure shape of the predatory Syrphidae communities found in orchard habitat and neighbouring shrubberies corroborated the values of  $H'$  general species diversity coefficient. It reached very similar values. The  $H'$  coefficient of the Syrphidae community in the orchards was 0.82. The  $H'$  coefficient of the Syrphidae community in the shrubberies was 0.76.

One definite eudominant in the orchard and shrubbery Syrphidae communities was confirmed with the values of  $J'$  uniformity index. It was identical for both habi-

tats: for the one in the orchard it reached 0.23 and for the shrubberies: 0.2.

A comparison of predatory Syrphidae communities in the orchard and neighbouring shrubberies in terms of quantity and the Re index proves that both communities were similar. The Re index reached a high value of 91%.

A quality comparison with the QS index shows that predatory Syrphidae communities in the orchard and shrubberies reached a 46.2% similarity in their species composition.

Only 6 predatory species were caught both in the orchard and shrubberies. They constituted 30% of the Syrphidae species composition and included the following species: *E. balteatus*, *E. corollae*, *M. mellinum*, *M. scalare*, *P. albimanus* and *S. pyrastris*.

In the orchard, 6 species only found in this habitat, namely: *D. venustus*, *E. eligans*, *M. auricollis*, *S. scripta*, *S. ribesii*, and *S. vitripennis*.

In the shrubberies, 8 species only found in this habitat, namely: *B. elongata*, *E. nitidicollis*, *L. lucorum*, *P. peltatus*, *P. scutatus*, *S. taeniata*, *S. torvus*, and *X. comtus*.

All the Syrphidae species fed on aphids (Aphidoidea) and Heteroptera larvae. *X. comtus* also feeds on Microlepidoptera larvae.




Fig. 1. The share of species dominant in the Syrphidae community in the orchard from 1999 to 2001




Fig. 2. The share of species dominant in the Syrphidae community caught in shrubberies from 1999 to 2001

## DISCUSSION AND CONCLUSIONS

The 1999–2001 studies of the Syrphidae infesting an orchard located in the Wielkopolska region of Poland, found a higher abundance of predators in the apple orchard than in those shrubberies in the vicinity. There was a higher abundance of Syrphidae in the orchard because the orchard was considerably larger than the area of the shrubberies. The orchard was overgrown with apple trees and consequently it had a higher abundance of aphids than the shrubberies. The presence of aphids as prey for Syrphidae larvae thus became a factor attracting Syrphidae imagines to the orchard. The abundance fluctuations in both habitats in particular years were similar, as was the quantity structure of the Syrphidae communities of the orchard and the shrubberies. A similar species diversity in the orchard and the shrubberies was reported and half the species composition was identical in both Syrphidae communities. A lower fluctuation in the number of species was found in the orchard than in the shrubberies. Thus, the results obtained prove that an apple orchard is a more attractive habitat for predatory Syrphidae than neighbouring shrubberies, and that predators can migrate between the orchard and the shrubberies. Wyss (1995) came to a similar conclusion when he stated that more aphid entomophages occurred on apple trees in the vicinity of wild plants than on apple trees without such plant life. This resulted in a lower number of aphids in the orchard neighbouring on wild plant life. Trzciński *et al.* (2011) found that in apple orchards neighbouring on abundant greenery in the form of shrubberies and a road overgrown with trees and bushes, there was a higher abundance of predatory Syrphidae than in the orchard neighbouring on agricultural cultivations. This can be explained by the appealing influence of the blooming wild plants for Syrphidae imagines, according to papers by Gilbert (1981), Wnuk and Gut (1994), Bostanian *et al.* (2004), Branquart and Hemptinne (2000), Colley and Luna (2000), Carreck and Williams (2002) and Ambrosino *et al.* (2006).

The research corroborated a decisive domination of *E. balteatus* in the orchard habitat, as this species had been previously reported by Wnuk (1972) as a dominant in that habitat. Also Trzciński *et al.* (2011) reported this species as an abundant one in Wielkopolska apple tree orchards. Miñarro and Dapena (2001) stated that *E. balteatus* was one of dominant predators of *Dysaphis plantaginea* (Pass.) in tree orchards in Spain. Weigel (1997) found that *E. balteatus* was a common species of apple orchards in Germany, while Hartfield (1997) reported it as such, in plum orchards in England.

Concluding, the orchard and neighbouring shrubberies in agroecosystems can act as alternative habitats for predatory Syrphidae. Such a conclusion can be drawn from similar fluctuations of the abundance of predatory Syrphidae in the orchard and shrubberies, a similar quantity structure of predators in both habitats, and a similar species diversity of entomophages. These results indicate the possibility of predators migrating from shrubberies to orchard and increasing the potential to control the abundance of Hemiptera species in the orchard. Thus, the shrubberies, due to their biocenotic functions, become a vital element of the agroecosystem structure.

## REFERENCES

- Ambrosino M.D., Luna J.M., Jepson P.C., Wratten S.D. 2006. Relative frequencies of visits to selected insectary plants by predatory hoverflies (Diptera: Syrphidae), other beneficial insects, and herbivores. *Environ. Entomol.* 35 (2): 394–400.
- Bennewicz J. 2011. Aphidivorous hoverflies (Diptera: Syrphidae) at field boundaries and woodland edges in an agricultural landscape. *Pol. Pismo Entomol.* 80 (1): 129–149.
- Bianchi F., Booij C.J.H., Tscharntke T. 2006. Sustainable pest regulation in agricultural landscapes: a review on landscape composition, biodiversity and natural pest control. *Proc. Royal Society B: Biol. Sci.* 273: 1715–1727.
- Bostanian N.J., Goulet H., O'Hara J., Masner L., Racette G. 2004. Towards insecticide free apple orchards: flowering plants to attract beneficial arthropods. *Biocontrol Sci. Technol.* 14 (1): 25–37.
- Branquart E., Hemptinne J.L. 2000. Selectivity in the exploitation of floral resources by hoverflies (Diptera: Syrphidae). *Ecography* 23: 732–742.
- Carreck N.L., Williams I.H. 2002. Food for insects pollinators on farmland: insect visits to flowers of annual seed mixtures. *J. Insect Conserv.* 6 (1): 13–23.
- Colley M.R., Luna J.M. 2000. Relative attractiveness of potential beneficial insectary plants to aphidophagous hoverflies (Diptera: Syrphidae). *Environ. Entomol.* 29 (5): 1054–1059.
- Debras J.F., Torre F., Rieux R., Reiter S., Garcin M.S., van Helden M., Buisson E., Dutoit T. 2006. Discrimination between agricultural management and the hedge effect in pear orchards (south-eastern France). *Ann. Appl. Biol.* 149 (3): 347–355.
- Gałecka B. 1995. Rola zadrzewień śródpolnych w funkcjonowaniu układu ofiara – drapieżca na przykładzie mszyc i afidofagów. *Zesz. Nauk. Komit. Człow. Śr.* 12: 99–111.
- Garbarkiewicz A., Trojanowski H. 1998. Występowanie mszycożernych *Syrphidae* na miedzach i przy drogach śródpolnych. [Aspects of balks and field – ways of conservation of Syrphidae in agriculture habitats]. *Prog. Plant Prot./Post. Ochr. Roślin* 38 (2): 631–623.
- Gilbert F.S. 1981. Foraging ecology of hoverflies: morphology of the mouthparts in relation to feeding on nectar and pollen in some common urban species. *Ecol. Entomol.* 6 (3): 245–262.
- Haenke S., Scheid B., Schaefer M., Tscharntke T., Thies C. 2009. Increasing syrphid fly diversity and density in sown flower strips within simple vs. complex landscape. *J. Appl. Ecol.* 46 (5): 1106–1114.
- Hartfield C.M. 1997. Biological control of aphid on plum. Ph.D. thesis. University of London, 196 pp.
- Kaczorowski G., Dębek-Jankowska A. 1997. Bzygowate (Syrphidae) różnych typów zarośli śródpolnych. [Hoverflies (Syrphidae) in different types of thickets]. *Prog. Plant Prot./Post. Ochr. Roślin* 37 (1): 85–87.
- Miñarro M., Dapena E. 2001. Predators of the rosy apple aphid, *Dysaphis plantaginea* (Pass.) in Asturian (NW Spain) apple orchards. *Integrated Fruit Production. IOBC/WPRS Bull.* 24 (5): 241–245.
- Moericke V. 1953. Wie finden geflügelte Blattläuse ihre Wirtspflanze? *Mitt. Biol. Reichsanst. Berlin* 75: 90–97.
- Pielou E.C. 1966. Shannon's formula as a measure of specific diversity; its use and misuse. *Am. Nat.* 100 (914): 463–465.

- Renkonen O. 1938. Statistisch-ökologische untersuchungen "über die terrestrische Käferwelt der finnischen Bruchmoore". Zool. Soc. Zoo. Bot. Fenn. 6: 1–226.
- Rossi J., Gamba U., Pinna M., Spagnolo S., Visentin C., Alma A. 2006. Hoverflies in organic apple orchards in north-western Italy. Bull. Insect. 59 (2): 111–114.
- Shannon C.E., Weaver W. 1963. The Mathematical Theory of Communication. Univ. Illinois Press, Urbana, 111 pp.
- Solomon M.G., Cross J.V., Fitzgerald J.D., Campbell C.A.M., Jolly R.L., Olszak R.W., Niemczyk E., Vogt H. 2000. Biocontrol of pests of apples and pears in northern and central Europe – 3. Predators. Biocontrol Sci. Techn. 10 (2): 91–128.
- Sørensen T. 1948. A method of establishing groups of equal amplitude in plant sociology based on similarity of species content and its application to analyses of vegetation on Danish common. Kong Danske Vidensk. Selsk Biol Skr. 5: 1–34.
- Trzciński P., Dolańska-Niedbała E., Piekarska-Boniecka H. 2011. Wpływ otoczenia sadu jabłoniowego na występowanie drapieżnego gatunku *Episyrphus balteatus* (Deg.) (Diptera, Syrphidae). [Effect of surroundings of apple orchard on the occurrence of predatory species *Episyrphus balteatus* (Deg.) (Diptera, Syrphidae)]. Prog. Plant Prot./Post. Ochr. Roślin 51 (4): 1792–1796.
- Trzciński P., Piekarska-Boniecka H. 2009. Występowanie bzyga prążkowanego *Episyrphus balteatus* (Deg.) (Diptera, Syrphidae) w sadach z integrowaną produkcją owoców i ich otoczeniu. [Occurrence of marmelade fly *Episyrphus balteatus* (Deg.) (Diptera, Syrphidae) in apple orchards with integrated production of fruit and their surroundings]. Prog. Plant Prot./Post. Ochr. Roślin 49 (4): 1987–1990.
- Weigel A. 1997. Zum Einfluß blühender Kräuter und Stauden in einer Apfelanlage auf die Populationsdynamik von Blattläusen und ihren Gegenspielern. Diplomarbeit, Universität Heidelberg, 123 pp.
- Wnuk A. 1972. Badania nad składem gatunkowym drapieżnych bzygowatych (Syrphidae, Diptera) występujących w koloniach mszyc na drzewach i krzewach owocowych. [Investigations on the species composition of predaceous Syrphidae (Diptera) occurring in the colonies of aphids on fruit trees and shrubs]. Pol. Pismo Entomol. 42 (1): 236–247.
- Wnuk A., Gut B. 1994. Atrakcyjność kwiatów dziko rosnących *Umbelliferae* dla mszycożernych *Syrphidae* (Dipt.). Pol. Pismo Entomol. 63 (1/2): 197–206.
- Wyss E. 1995. The effects of weed strips on aphids and aphidophagous predators in an apple orchard. Entomol. Exp. Appl. 75 (1): 43–49.